

RHOC Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant RHOC. Catalog # AT3636a

Product Information

ApplicationWB, EPrimary AccessionP08134Other AccessionBC007245

Reactivity Human, Mouse, Rat

Host mouse
Clonality monoclonal
Isotype IgG2b Kappa
Clone Names 2.0E+12
Calculated MW 22006

Additional Information

Gene ID 389

Other Names Rho-related GTP-binding protein RhoC, Rho cDNA clone 9, h9, RHOC, ARH9,

ARHC

Target/Specificity RHOC (AAH07245, 1 a.a. ~ 193 a.a) full-length recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions RHOC Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.


Background

This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The protein encoded by this gene is prenylated at its C-terminus, and localizes to the cytoplasm and plasma membrane. It is thought to be important in cell locomotion. Overexpression of this gene is associated with tumor cell proliferation and metastasis. Multiple alternatively spliced variants, encoding the same protein, have been identified.


References

1.Epidermal Growth Factor Stimulates Human Trophoblast Cell Migration through Rho A and Rho C Activation.Han J, Li L, Hu J, Yu L, Zheng Y, Guo J, Zheng X, Yi P, Zhou Y.Endocrinology. 2010 Feb 11. [Epub ahead of print]

Images




Western Blot analysis of RHOC expression in transfected



293T cell line by RHOC monoclonal antibody (M01), clone 2E12.

Lane 1: RHOC transfected lysate(22 KDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged RHOC is approximately 3ng/ml as a capture antibody.

Citations

• Knockdown of RHOC by shRNA suppresses invasion and migration of cholangiocellular carcinoma cells via inhibition of MMP2, MMP3, MMP9 and epithelial-mesenchymal transition.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.