

# RPL14 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant RPL14. Catalog # AT3703a

#### **Product Information**

Application WB, IHC, E
Primary Accession P50914
Other Accession NM\_003973
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2a Kappa

Clone Names 1B4 Calculated MW 23432

### **Additional Information**

**Gene ID** 9045

Other Names 60S ribosomal protein L14, CAG-ISL 7, RPL14

**Target/Specificity** RPL14 (NP\_003964, 1 a.a. ~ 90 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IHC~~1:100~500 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** RPL14 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.


## **Background**

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L14E family of ribosomal proteins. It contains a basic region-leucine zipper (bZIP)-like domain. The protein is located in the cytoplasm. This gene contains a trinucleotide (GCT) repeat tract whose length is highly polymorphic; these triplet repeats result in a stretch of alanine residues in the encoded protein. Transcript variants utilizing alternative polyA signals and alternative 5'-terminal exons exist but all encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.


### References

The role of human ribosomal proteins in the maturation of rRNA and ribosome production. Robledo S, et al. RNA, 2008 Sep. PMID 18697920. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. Olsen JV, et al. Cell, 2006 Nov 3. PMID 17081983. Nucleolar proteome dynamics. Andersen JS, et al. Nature, 2005 Jan 6. PMID 15635413. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.


### **Images**



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (35.64 KDa) .



Immunoperoxidase of monoclonal antibody to RPL14 on formalin-fixed paraffin-embedded human colon. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged RPL14 is approximately 3ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.