

# SGK2 Antibody (monoclonal) (M07)

Mouse monoclonal antibody raised against a partial recombinant SGK2. Catalog # AT3856a

### **Product Information**

**Application** WB, IF, E **Primary Accession** Q9HBY8 **Other Accession** BC065511 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 Kappa **Clone Names** 2F6

#### **Additional Information**

**Calculated MW** 

**Gene ID** 10110

**Other Names** Serine/threonine-protein kinase Sgk2, Serum/glucocorticoid-regulated kinase

2, SGK2

41175

**Target/Specificity** SGK2 (AAH65511, 293 a.a. ~ 367 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IF~~1:50~200 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** SGK2 Antibody (monoclonal) (M07) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Background**


This gene encodes a serine/threonine protein kinase. Although this gene product is similar to serum- and glucocorticoid-induced protein kinase (SGK), this gene is not induced by serum or glucocorticoids. This gene is induced in response to signals that activate phosphatidylinositol 3-kinase, which is also true for SGK. Two alternate transcripts encoding two different isoforms have been described.

### References


Sgk kinases and their role in epithelial transport. Loffing J, et al. Annu Rev Physiol, 2006. PMID 16460280. Transcriptional maps of 10 human chromosomes at 5-nucleotide resolution. Cheng J, et al.

Science, 2005 May 20. PMID 15790807.Regulation of cardiac shal-related potassium channel Kv 4.3 by serum- and glucocorticoid-inducible kinase isoforms in Xenopus oocytes. Baltaev R, et al. Pflugers Arch, 2005 Apr. PMID 15578212.The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.Regulation of KCNE1-dependent K(+) current by the serum and glucocorticoid-inducible kinase (SGK) isoforms. Embark HM, et al. Pflugers Arch, 2003 Feb. PMID 12634932.


### **Images**




Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (33.88 KDa) .



SGK2 monoclonal antibody (M07), clone 2F6 Western Blot analysis of SGK2 expression in IMR-32 ( (Cat # AT3856a )



Immunofluorescence of monoclonal antibody to SGK2 on HeLa cell. [antibody concentration 30 ug/ml]



Detection limit for recombinant GST tagged SGK2 is approximately 0.03ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.