

TLK2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant TLK2. Catalog # AT4251a

Product Information

Application WB, IHC, IF **Primary Accession Q86UE8 Other Accession** BC044925 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 kappa **Clone Names** 2G7 Calculated MW 87661

Additional Information

Gene ID 11011

Other Names Serine/threonine-protein kinase tousled-like 2, HsHPK, PKU-alpha,

Tousled-like kinase 2, TLK2

Target/Specificity TLK2 (AAH44925, 1 a.a. ~ 126 a.a) partial recombinant protein with GST tag.

MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions TLK2 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.

Background


The Tousled-like kinases, first described in Arabidopsis, are nuclear serine/threonine kinases that are potentially involved in the regulation of chromatin assembly.

References


Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Association of mitotic regulation pathway polymorphisms with pancreatic cancer risk and outcome. Couch FJ, et al. Cancer Epidemiol Biomarkers Prev, 2010 Jan. PMID 20056645. Phosphorylation-mediated control of histone chaperone ASF1 levels by

Tousled-like kinases. Pilyugin M, et al. PLoS One, 2009 Dec 16. PMID 20016786.Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931.FEZ1 dimerization and interaction with transcription regulatory proteins involves its coiled-coil region. Assmann EM, et al. J Biol Chem, 2006 Apr 14. PMID 16484223.


Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (39.27 KDa) .



Immunoperoxidase of monoclonal antibody to TLK2 on formalin-fixed paraffin-embedded human colon tissue. [antibody concentration 5 ug/ml]



Immunofluorescence of monoclonal antibody to TLK2 on HeLa cell. [antibody concentration 10 ug/ml]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.