

# TLX3 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant TLX3. Catalog # AT4260a

### **Product Information**

**Application** WB, E **Primary Accession** 043711 Other Accession NM 021025 Reactivity Human Host mouse Clonality monoclonal Isotype IgG1 Kappa **Clone Names** 2A3 Calculated MW 31867

#### **Additional Information**

**Gene ID** 30012

Other Names T-cell leukemia homeobox protein 3, Homeobox protein Hox-11L2, TLX3,

HOX11L2

Target/Specificity TLX3 (NP\_066305, 192 a.a. ~ 291 a.a) partial recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** TLX3 Antibody (monoclonal) (M01) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Background**


RNX (HOX11L2, TLX3) belongs to a family of orphan homeobox genes that encode DNA-binding nuclear transcription factors. Members of the HOX11 gene family are characterized by a threonine-47 replacing cytosine in the highly conserved homeodomain (Dear et al., 1993 [PubMed 8099440]).

### References


Thymic adult T-cell acute lymphoblastic leukemia stratified in standard- and high-risk group by aberrant HOX11L2 expression: experience of the German multicenter ALL study group. Baak U, et al. Leukemia, 2008 Jun. PMID 18368072.Cooperative genetic defects in TLX3 rearranged pediatric T-ALL. Van Vlierberghe P, et al.

Leukemia, 2008 Apr. PMID 18185524. Activation of TLX3 and NKX2-5 in t(5;14)(q35;q32) T-cell acute lymphoblastic leukemia by remote 3'-BCL11B enhancers and coregulation by PU.1 and HMGA1. Nagel S, et al. Cancer Res, 2007 Feb 15. PMID 17308084. Nuclear factor Y drives basal transcription of the human TLX3, a gene overexpressed in T-cell acute lymphocytic leukemia. Borghini S, et al. Mol Cancer Res, 2006 Sep. PMID 16966433. Significance of HOX11L2/TLX3 expression in children with T-cell acute lymphoblastic leukemia treated on Children's Cancer Group protocols. Gottardo NG, et al. Leukemia, 2005 Sep. PMID 15990867.


### **Images**



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa).




TLX3 monoclonal antibody (M01), clone 2A3 Western Blot analysis of TLX3 expression in MCF-7 ( (Cat # AT4260a )




Western Blot analysis of TLX3 expression in transfected 293T cell line by TLX3 monoclonal antibody (M01), clone 2A3.

Lane 1: TLX3 transfected lysate(31.9 KDa). Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged TLX3 is approximately 0.03ng/ml as a capture antibody.

Western blot analysis of TLX3 over-expressed 293 cell



line, cotransfected with TLX3 Validated Chimera RNAi ( (Cat # AT4260a )

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.