

TTF2 Antibody (monoclonal) (M12)

Mouse monoclonal antibody raised against a full length recombinant TTF2. Catalog # AT4390a

Product Information

Application WB, E
Primary Accession Q9UNY4
Other Accession NM_003594
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2a Kappa

Clone Names 2B6 Calculated MW 129588

Additional Information

Gene ID 8458

Other Names Transcription termination factor 2, 364-, Lodestar homolog, RNA polymerase

II termination factor, Transcription release factor 2, F2, HuF2, TTF2

Target/Specificity TTF2 (NP_003585, 385 a.a. ~ 509 a.a) full length recombinant protein with GST

tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions TTF2 Antibody (monoclonal) (M12) is for research use only and not for use in

diagnostic or therapeutic procedures.

Background


This gene encodes a member of the SWI2/SNF2 family of proteins, which play a critical role in altering protein-DNA interactions. The encoded protein has been shown to have dsDNA-dependent ATPase activity and RNA polymerase II termination activity. This protein interacts with cell division cycle 5-like, associates with human splicing complexes, and plays a role in pre-mRNA splicing.

References

Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. The interactome of the histone gene regulatory factor HiNF-P suggests novel cell cycle


related roles in transcriptional control and RNA processing. Miele A, et al. J Cell Biochem, 2007 Sep 1. PMID 17577209. The DNA sequence and biological annotation of human chromosome 1. Gregory SG, et al. Nature, 2006 May 18. PMID 16710414. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Rescue of the TTF2 knockdown phenotype with an siRNA-resistant replacement vector. Jiang Y, et al. Cell Cycle, 2004 Sep. PMID 15467445.

Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (39.75 KDa) .

TTF2 monoclonal antibody (M12), clone 2B6 Western Blot analysis of TTF2 expression in MES-SA/Dx5 ((Cat # AT4390a)



Detection limit for recombinant GST tagged TTF2 is approximately 1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.