

# TWIST1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant TWIST1.

Catalog # AT4410a

## Product Information

---

<b>Application</b>	WB, IP, E
<b>Primary Accession</b>	<a href="#">Q15672</a>
<b>Other Accession</b>	<a href="#">NM_000474</a>
<b>Reactivity</b>	Human, Rat
<b>Host</b>	Mouse
<b>Clonality</b>	monoclonal
<b>Isotype</b>	IgG1 Kappa
<b>Clone Names</b>	3E12
<b>Calculated MW</b>	20954

## Additional Information

---

<b>Gene ID</b>	7291
<b>Other Names</b>	Twist-related protein 1, Class A basic helix-loop-helix protein 38, bHLHa38, H-twist, TWIST1, BHLHA38, TWIST
<b>Target/Specificity</b>	TWIST1 (NP_000465, 100 a.a. ~ 202 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
<b>Dilution</b>	WB~~1:500~1000 IP~~N/A E~~N/A
<b>Format</b>	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
<b>Storage</b>	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
<b>Precautions</b>	TWIST1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

## Background

---

Basic helix-loop-helix (bHLH) transcription factors have been implicated in cell lineage determination and differentiation. The protein encoded by this gene is a bHLH transcription factor and shares similarity with another bHLH transcription factor, Dermo1. The strongest expression of this mRNA is in placental tissue; in adults, mesodermally derived tissues express this mRNA preferentially. Mutations in this gene have been found in patients with Saethre-Chotzen syndrome.


## References

---


- 1.Cellular migration and invasion uncoupled: Increased migration is not an inexorable consequence of

EMT.Schaeffer D, Somarelli JA, Hanna G, Palmer GM, Garcia-Blanco MAMol Cell Biol. 2014 Jul 7. pii: MCB.00694-14.2.EMT and induction of miR-21 mediate metastasis development in Trp53-deficient tumours.Bornachea O, Santos M, Martinez-Cruz AB, Garcia-Escudero R, Duenas M, Costa C, Segrelles C, Lorz C, Buitrago A, Saiz-Ladera C, Agirre X, Grande T, Paradela B, Maraver A, Ariza JM, Prosper F, Serrano M, Sanchez-Cespedes M, Paramio JM.Sci Rep. 2012;2:434. Epub 2012 May 31.3.Conversion of Helicobacter pylori CagA from senescence inducer to oncogenic driver through polarity-dependent regulation of p21.Saito Y, Murata-Kamiya N, Hirayama T, Ohba Y, Hatakeyama M.J Exp Med. 2010 Sep 27;207(10):2157-74. Epub 2010 Sep 20.


## Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.07 KDa) .




TWIST1 monoclonal antibody (M01), clone 3E11. Western Blot analysis of TWIST1 expression in rat brain.




Western Blot analysis of TWIST1 expression in transfected 293T cell line by TWIST1 monoclonal antibody (M01), clone 3E11.

Lane 1: TWIST1 transfected lysate(21 KDa).  
Lane 2: Non-transfected lysate.

Immunoprecipitation of TWIST1 transfected lysate using anti-TWIST1 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with TWIST1 MaxPab rabbit polyclonal antibody.



Detection limit for recombinant GST tagged TWIST1 is approximately 3ng/ml as a capture antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.