

# TXNDC4 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant TXNDC4. Catalog # AT4417a

#### **Product Information**

Application WB, IHC, IP, E
Primary Accession Q9BS26
Other Accession BC005374
Reactivity Human
Host mouse
Clonality monoclonal
Isotype IgG2b Kappa

Clone Names 3C7
Calculated MW 46971

### **Additional Information**

**Gene ID** 23071

Other Names Endoplasmic reticulum resident protein 44, ER protein 44, ERp44, Thioredoxin

domain-containing protein 4, ERP44, KIAA0573, TXNDC4

**Target/Specificity** TXNDC4 (AAH05374, 30 a.a. ~ 406 a.a) full-length recombinant protein with

GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 IHC~~1:100~500 IP~~N/A E~~N/A

**Format** Clear, colorless solution in phosphate buffered saline, pH 7.2.

**Storage** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.


**Precautions** TXNDC4 Antibody (monoclonal) (M01) is for research use only and not for use

in diagnostic or therapeutic procedures.


### References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.Crystal structure of human ERp44 shows a dynamic functional modulation by its carboxy-terminal tail. Wang L, et al. EMBO Rep, 2008 Jul. PMID 18552768.Multistep, sequential control of the trafficking and function of the multiple sulfatase deficiency gene product, SUMF1 by PDI, ERGIC-53 and ERp44. Fraldi A, et al. Hum Mol Genet, 2008 Sep 1. PMID 18508857.

## **Images**




Immunoperoxidase of monoclonal antibody to TXNDC4 on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 3 ug/ml]





Immunoprecipitation of TXNDC4 transfected lysate using anti-TXNDC4 monoclonal antibody and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with TXNDC4 MaxPab rabbit polyclonal antibody.



Detection limit for recombinant GST tagged TXNDC4 is 1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.