

UBTF Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant UBTF.

Catalog # AT4454a

Product Information

Application	WB, IHC, IF, E
Primary Accession	P17480
Other Accession	NM_014233
Reactivity	Human, Rat
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	1A2
Calculated MW	89406

Additional Information

Gene ID	7343
Other Names	Nucleolar transcription factor 1, Autoantigen NOR-90, Upstream-binding factor 1, UBF-1, UBTF, UBF, UBF1
Target/Specificity	UBTF (NP_055048, 551 a.a. ~ 650 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	UBTF Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

Upstream binding factor (UBF) is a transcription factor required for expression of the 18S, 5.8S, and 28S ribosomal RNAs, along with SL1 (a complex of TBP (MIM 600075) and multiple TBP-associated factors or 'TAFs'). Two UBF polypeptides, of 94 and 97 kD, exist in the human (Bell et al., 1988 [PubMed 3413483]). UBF is a nucleolar phosphoprotein with both DNA binding and transactivation domains. Sequence-specific DNA binding to the core and upstream control elements of the human rRNA promoter is mediated through several HMG boxes (Jantzen et al., 1990 [PubMed 2330041]).


References

Molecular mechanisms underlying nucleocytoplasmic shuttling of actinin-4. Kumeta M, et al. J Cell Sci, 2010 Apr 1. PMID 20197409. Transcriptional corepressor TLE1 functions with Runx2 in epigenetic repression of ribosomal RNA genes. Ali SA, et al. Proc Natl Acad Sci U S A, 2010 Mar 2. PMID 20160071. Treacle recruits RNA polymerase I complex to the nucleolus that is independent of UBF. Lin CI, et al. Biochem Biophys Res Commun, 2009 Aug 21. PMID 19527688. Temperature-induced partially unfolded state of hUBF HMG Box-5: conformational and dynamic investigations of the Box-5 thermal intermediate ensemble. Liu Z, et al. Proteins, 2009 Nov 1. PMID 19452555. Nucleolar localization of upstream binding factor in HeLa cells depends on rRNA synthesis activities. He J, et al. Folia Biol (Praha), 2008. PMID 19393134.


Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .




UBTF monoclonal antibody (M03), clone 1A2. Western Blot analysis of UBTF expression in PC-12 (Cat # AT4454a)




UBTF monoclonal antibody (M03), clone 1A2 Western Blot analysis of UBTF expression in HepG2 (Cat # AT4454a)



Immunoperoxidase of monoclonal antibody to UBTF on formalin-fixed paraffin-embedded human placenta. [antibody concentration 0.5 ug/ml]



Immunofluorescence of monoclonal antibody to UBTF on HepG2 cell. [antibody concentration 10 ug/ml]



Detection limit for recombinant GST tagged UBTF is approximately 1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.