

USP52 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant USP52.

Catalog # AT4496a

Product Information

Application	WB, IF
Primary Accession	Q504Q3
Other Accession	NM_014871
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	6A7
Calculated MW	135368

Additional Information

Gene ID	9924
Other Names	PAB-dependent poly(A)-specific ribonuclease subunit PAN2 {ECO:0000255 HAMAP-Rule:MF_03182}, hPan2, 31134 {ECO:0000255 HAMAP-Rule:MF_03182}, Inactive ubiquitin carboxyl-terminal hydrolase 52 {ECO:0000255 HAMAP-Rule:MF_03182}, PAB1P-dependent poly(A)-nuclease {ECO:0000255 HAMAP-Rule:MF_03182}, PAN deadenylation complex catalytic subunit 2 {ECO:0000255 HAMAP-Rule:MF_03182}, PAN2 {ECO:0000255 HAMAP-Rule:MF_03182}, KIAA0710, USP52
Target/Specificity	USP52 (NP_055686, 1099 a.a. ~ 1198 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IF~~1:50~200
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	USP52 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


Background

This gene encodes a deadenylase that functions as the catalytic subunit of the polyadenylate binding protein dependent poly(A) nuclease complex. The encoded protein is a magnesium dependent 3' to 5' exoribonuclease that is involved in the degradation of cytoplasmic mRNAs. Alternate splicing results in multiple transcript variants.


References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. Mechanism of mRNA deadenylation: evidence for a molecular interplay between translation termination factor eRF3 and mRNA deadenylases. Funakoshi Y, et al. Genes Dev, 2007 Dec 1. PMID 18056425. Human TOB, an antiproliferative transcription factor, is a poly(A)-binding protein-dependent positive regulator of cytoplasmic mRNA deadenylation. Ezzeddine N, et al. Mol Cell Biol, 2007 Nov. PMID 17785442.

Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Immunofluorescence of monoclonal antibody to PAN2 on HeLa cell . [antibody concentration 10 ug/ml]

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.