

ZBTB16 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ZBTB16.

Catalog # AT4569a

Product Information

Application	WB, E
Primary Accession	Q05516
Other Accession	BC029812
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2a Kappa
Clone Names	3A7
Calculated MW	74274

Additional Information

Gene ID	7704
Other Names	Zinc finger and BTB domain-containing protein 16, Promyelocytic leukemia zinc finger protein, Zinc finger protein 145, Zinc finger protein PLZF, ZBTB16, PLZF, ZNF145
Target/Specificity	ZBTB16 (AAH29812, 381 a.a. ~ 480 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	ZBTB16 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.




Background

This gene is a member of the Krueppel C2H2-type zinc-finger protein family and encodes a zinc finger transcription factor that contains nine Kruppel-type zinc finger domains at the carboxyl terminus. This protein is located in the nucleus, is involved in cell cycle progression, and interacts with a histone deacetylase. Specific instances of aberrant gene rearrangement at this locus have been associated with acute promyelocytic leukemia (APL). Alternate transcriptional splice variants have been characterized.

References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Application of gene network analysis techniques identifies AXIN1/PDIA2 and endoglin haplotypes associated with bicuspid aortic valve. Wooten EC, et al. PLoS One, 2010 Jan 21. PMID 20098615. Generation of PLZF+ CD4+ T cells via MHC class II-dependent thymocyte-thymocyte interaction is a physiological process in humans. Lee YJ, et al. J Exp Med, 2010 Jan 18. PMID 20038602. The promyelocytic leukemia zinc-finger gene, PLZF, is frequently downregulated in malignant mesothelioma cells and contributes to cell survival. Cheung M, et al. Oncogene, 2010 Mar 18. PMID 20010871.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.