

ZIC1 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant ZIC1.

Catalog # AT4579a

Product Information

Application	WB, E
Primary Accession	Q15915
Other Accession	NM_003412
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG2b Kappa
Clone Names	4D11
Calculated MW	48309

Additional Information

Gene ID	7545
Other Names	Zinc finger protein ZIC 1, Zinc finger protein 201, Zinc finger protein of the cerebellum 1, ZIC1, ZIC, ZNF201
Target/Specificity	ZIC1 (NP_003403, 2 a.a. ~ 95 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	ZIC1 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

Background


This gene encodes a member of the ZIC family of C2H2-type zinc finger proteins. Members of this family are important during development. Aberrant expression of this gene is seen in medulloblastoma, a childhood brain tumor. This gene is closely linked to the gene encoding zinc finger protein of the cerebellum 4, a related family member on chromosome 3. This gene encodes a transcription factor that can bind and transactivate the apolipoprotein E gene.

References


ZIC1 is downregulated through promoter hypermethylation in gastric cancer. Wang LJ, et al. Biochem

Biophys Res Commun, 2009 Feb 20. PMID 19135984. Genome-wide association analysis of susceptibility and clinical phenotype in multiple sclerosis. Baranzini SE, et al. Hum Mol Genet, 2009 Feb 15. PMID 19010793. Inhibition of translation in living eukaryotic cells by an RNA G-quadruplex motif. Arora A, et al. RNA, 2008 Jul. PMID 18515550. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. ZIC1 gene expression is controlled by DNA and histone methylation in mesenchymal proliferations. Pourebrahim R, et al. FEBS Lett, 2007 Oct 30. PMID 17936758.

Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.08 kDa) .



Detection limit for recombinant GST tagged ZIC1 is approximately 1 ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.