

ZNF92 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant ZNF92.

Catalog # AT4646a

Product Information


Application	WB, IHC, E
Primary Accession	Q03936
Other Accession	NM_152626
Reactivity	Human
Host	mouse
Clonality	monoclonal
Isotype	IgG1 Kappa
Clone Names	1F2
Calculated MW	68487

Additional Information


Gene ID	168374
Other Names	Zinc finger protein 92, Zinc finger protein HTF12, ZNF92
Target/Specificity	ZNF92 (NP_689839, 490 a.a. ~ 586 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.
Dilution	WB~~1:500~1000 IHC~~1:100~500 E~~N/A
Format	Clear, colorless solution in phosphate buffered saline, pH 7.2 .
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.
Precautions	ZNF92 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.41 KDa) .




Western Blot analysis of ZNF92 expression in transfected 293T cell line by ZNF92 monoclonal antibody (M01), clone 1F2.

Lane 1: ZNF92 transfected lysate(60.2 KDa).
Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to ZNF92 on formalin-fixed paraffin-embedded human prostate. [antibody concentration 1.2 ug/ml]



Detection limit for recombinant GST tagged ZNF92 is approximately 0.1ng/ml as a capture antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.