10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

USP11 Antibody (N-term)

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AW5070

Product Information

Application	FC, WB
Primary Accession	P51784
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	109817
Isotype	IgG1,κ
Antigen Source	HUMAN

Additional Information

Antigen Region	32-300
Other Names	Ubiquitin carboxyl-terminal hydrolase 11, Deubiquitinating enzyme 11, Ubiquitin thioesterase 11, Ubiquitin-specific-processing protease 11, USP11, UHX1
Dilution	FC~~1:25 WB~~1:1000
Target/Specificity	This USP11 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 32-300 amino acids from the N-terminal region of human USP11.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	USP11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	USP11
Synonyms	UHX1
Function	Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains (PubMed:12084015, PubMed:15314155,

PubMed:17897950, PubMed:19874889, PubMed:20233726, PubMed:24724799, PubMed:28992046). Inhibits the degradation of target proteins by the proteasome (PubMed:12084015). Cleaves preferentially 'Lys-6' and 'Lys- 63'-linked ubiquitin chains. Has lower activity with 'Lys-11' and 'Lys-33'-linked ubiquitin chains, and extremely low activity with 'Lys-27', 'Lys-29' and 'Lys-48'-linked ubiquitin chains (in vitro) (PubMed:24724799). Plays a role in the regulation of pathways leading to NF-kappa-B activation (PubMed:17897950, PubMed:19874889). Plays a role in the regulation of DNA repair after double-stranded DNA breaks (PubMed:15314155, PubMed:20233726). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:20601937). Promotes cell proliferation by deubiquitinating phosphorylated E2F1 (PubMed:28992046).

Cellular Location

Nucleus. Cytoplasm. Chromosome. Note=Predominantly nuclear (PubMed:12084015, PubMed:15314155). Associates with chromatin (PubMed:20233726, PubMed:20601937).


Background

Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains. Inhibits the degradation of target proteins by the proteasome. Plays a role in the regulation of pathways leading to NF-kappa-B activation. Plays a role in the regulation of DNA repair after double-stranded DNA breaks.

References


Ross M.T.,et al.Nature 434:325-337(2005).
Ideguchi H.,et al.Biochem. J. 367:87-95(2002).
Swanson D.A.,et al.Hum. Mol. Genet. 5:533-538(1996).
Schoenfeld A.R.,et al.Mol. Cell. Biol. 24:7444-7455(2004).
Yamaguchi T.,et al.J. Biol. Chem. 282:33943-33948(2007).

Images



Western blot analysis of lysates from Jurkat,Hela cell line (from left to right), using USP11 Antibody (C-term R565)(Cat. #AW5070). AW5070 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

Flow cytometric analysis of Hela cells using USP11 Antibody (C-term R565)(green, Cat#AW5070) compared to an isotype control of mouse IgG1(blue). AW5070 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.