

# USP5 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AW5077

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P45974</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Calculated MW</b>	95786
<b>Isotype</b>	IgG1, $\kappa$
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	8078
<b>Antigen Region</b>	1-160
<b>Other Names</b>	Ubiquitin carboxyl-terminal hydrolase 5, Deubiquitinating enzyme 5, Isopeptidase T, Ubiquitin thioesterase 5, Ubiquitin-specific-processing protease 5, USP5, ISOT
<b>Dilution</b>	WB~~1:1000
<b>Target/Specificity</b>	This USP5 antibody is generated from a mouse immunized with a recombination protein from the human region of human USP5.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	USP5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	USP5
<b>Synonyms</b>	ISOT
<b>Function</b>	Deubiquitinating enzyme that participates in a wide range of cellular

processes by specifically cleaving isopeptide bonds between ubiquitin and substrate proteins or ubiquitin itself. Affects thereby important cellular signaling pathways such as NF-kappa-B, Wnt/beta- catenin, and cytokine production by regulating ubiquitin-dependent protein degradation. Participates in the activation of the Wnt signaling pathway by promoting FOXM1 deubiquitination and stabilization that induces the recruitment of beta-catenin to Wnt target gene promoter (PubMed:[26912724](#)). Regulates the assembly and disassembly of heat-induced stress granules by mediating the hydrolysis of unanchored ubiquitin chains (PubMed:[29567855](#)). Promotes lipopolysaccharide-induced apoptosis and inflammatory response by stabilizing the TXNIP protein (PubMed:[37534934](#)). Affects T-cell biology by stabilizing the inhibitory receptor on T-cells PDC1 (PubMed:[37208329](#)). Acts as a negative regulator of autophagy by regulating ULK1 at both protein and mRNA levels (PubMed:[37607937](#)). Acts also as a negative regulator of type I interferon production by simultaneously removing both 'Lys-48'-linked unanchored and 'Lys-63'-linked anchored polyubiquitin chains on the transcription factor IRF3 (PubMed:[39761299](#)). Modulates the stability of DNA mismatch repair protein MLH1 and counteracts the effect of the ubiquitin ligase UBR4 (PubMed:[39032648](#)). Upon activation by insulin, it gets phosphorylated through mTORC1-mediated phosphorylation to enhance YTHDF1 stability by removing 'Lys-11'-linked polyubiquitination (PubMed:[39900921](#)). May also deubiquitinate other substrates such as the calcium channel CACNA1H (By similarity).

#### Cellular Location

Cytoplasm. Cytoplasm, Stress granule. Nucleus


## Background

Cleaves linear and branched multiubiquitin polymers with a marked preference for branched polymers. Involved in unanchored 'Lys-48'-linked polyubiquitin disassembly. Binds linear and 'Lys-63'-linked polyubiquitin with a lower affinity. Knock-down of USP5 causes the accumulation of p53/TP53 and an increase in p53/TP53 transcriptional activity because the unanchored polyubiquitin that accumulates is able to compete with ubiquitinated p53/TP53 but not with MDM2 for proteasomal recognition.

## References

- Falquet L.,et al.FEBS Lett. 376:233-237(1995).  
 Ansari-Lari M.A.,et al.Genome Res. 6:314-326(1996).  
 Ansari-Lari M.A.,et al.Genome Res. 7:268-280(1997).  
 Tashayev V.L.,et al.Submitted (NOV-1995) to the EMBL/GenBank/DDBJ databases.  
 Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

## Images



Western blot analysis of lysates from mouse brain tissue, PC-3 cell line, rat liver tissue, U-251 MG cell line, rat brain tissue (from left to right), using USP5 Antibody (Cat. #AW5077). AW5077 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.