

ADRA1D Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5131

Product Information

Application	IHC-P, FC, WB
Primary Accession	P25100
Reactivity	Human, Rat
Predicted	Rabbit, Dog, Sheep, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60463
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	146
Antigen Region	1-30
Other Names	Alpha-1D adrenergic receptor, Alpha-1A adrenergic receptor, Alpha-1D adrenoceptor, Alpha-1D adrenoceptor, Alpha-adrenergic receptor 1a, ADRA1D, ADRA1A
Dilution	WB~~1:1000 FC~~1:25 IHC-P~~1:100~500
Target/Specificity	This ADRA1D antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1-30amino acids from the N-terminal region of human ADRA1D.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ADRA1D Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ADRA1D
-------------	--------

Synonyms	ADRA1A
Function	This alpha-adrenergic receptor mediates its effect through the influx of extracellular calcium.
Cellular Location	Cell membrane; Multi-pass membrane protein.


Background

This alpha-adrenergic receptor mediates its effect through the influx of extracellular calcium.


References

- Bruno J.F.,et al.Biochem. Biophys. Res. Commun. 179:1485-1490(1991).
 Forray C.,et al.Mol. Pharmacol. 45:703-708(1994).
 Schwinn D.A.,et al.J. Pharmacol. Exp. Ther. 272:134-142(1995).
 Weinberg D.H.,et al.Biochem. Biophys. Res. Commun. 201:1296-1304(1994).
 Esbenshade T.A.,et al.Mol. Pharmacol. 47:977-985(1995).

Images




Western blot analysis of lysates from A549 cell line, rat brain tissue, LNCaP, PC-3 cell line (from left to right), using ADRA1D Antibody (N-term)(Cat. #AW5131). AW5131 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.




Flow cytometric analysis of MCF-7 cells using ADRA1D Antibody (N-term)(green, Cat#AW5131) compared to an isotype control of rabbit IgG(blue). AW5131 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

Immunohistochemical analysis of paraffin-embedded H. kidney section using ADRA1D Antibody (N-term)(Cat#AW5131). AW5131 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. small intestine section using ADRA1D Antibody (N-term)(Cat#AW5131). AW5131 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded R. kidney section using ADRA1D Antibody (N-term)(Cat#AW5131). AW5131 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

