

Phospho-HIST1H3B3(S10) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5153

Product Information

Application	WB
Primary Accession	P68431
Reactivity	Human
Predicted	Mouse, Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15404
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	8350;8351;8352;8353;8354;8355;8356;8357;8358;8968
Antigen Region	8-35
Other Names	HIST1H3A; H3FA; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; Histone H3/h; Histone H3/i; Histone H3/j; Histone H3/k; Histone H3/l
Dilution	WB~~1:500
Target/Specificity	This HIST1H3B3 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S10 of human HIST1H3B3.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-HIST1H3B3(S10) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	H3C1 (HGNC:4766)
Synonyms	H3FA, HIST1H3A

Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.


Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.


References

Lusic, M., et al., EMBO J. 22(24):6550-6561 (2003).
Deng, L., et al., Virology 289(2):312-326 (2001).
Deng, L., et al., Virology 277(2):278-295 (2000).
El Kharroubi, A., et al., Mol. Cell. Biol. 18(5):2535-2544 (1998).
Albig, W., et al., Hum. Genet. 101(3):284-294 (1997).

Images



Western blot analysis of lysate from HeLa cell line, using Phospho-H3-pS10 Antibody (Cat. #AW5153). AW5153 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.



Western blot analysis of lysate from HeLa cell line, using Phospho-H3-pS10 Antibody (Cat. #AW5153). AW5153 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.