

Phospho-M ERBB2(Y1140) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5161

Product Information

Application	WB
Primary Accession	P70424
Other Accession	NP_001003817.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	138579
Isotype	Rabbit IgG
Antigen Source	MOUSE

Additional Information

Gene ID	13866
Antigen Region	1125-1147
Other Names	ErbB2; Kias3023; Neu; Receptor tyrosine-protein kinase erbB-2; Proto-oncogene Neu; Proto-oncogene c-ErbB-2; p185erbB2; CD_antigen=CD340; Flags: Precursor
Dilution	WB~~1:1000
Target/Specificity	This mouse ERBB2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y1140 of mouse ERBB2.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Phospho-M ERBB2(Y1140) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ErbB2
Synonyms	Kias3023, Neu

Function	Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization (By similarity).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P04626}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P04626} Cell projection, ruffle membrane {ECO:0000250 UniProtKB:P04626}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P04626} Early endosome {ECO:0000250 UniProtKB:P04626}. Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:P04626}. Nucleus {ECO:0000250 UniProtKB:P04626}. Note=Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1. Also detected in endosome-to-TGN retrograde vesicles. Internalized from the cell membrane in response to EGF stimulation. {ECO:0000250 UniProtKB:P04626}
Tissue Location	Expressed predominantly in uterine epithelial cells. In the muscle, expression localizes to the synaptic sites of muscle fibers

Background


This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized.

References

Cabodi, S., et al. FASEB J. 24(10):3796-3808(2010)
 Johnson, E., et al. J. Biol. Chem. 285(38):29491-29501(2010)
 Huck, L., et al. Proc. Natl. Acad. Sci. U.S.A. 107(35):15559-15564(2010)
 Chuang, T.D., et al. J. Biol. Chem. 285(31):23598-23606(2010)
 Simeone, L., et al. J. Neurosci. 30(19):6620-6634(2010)

Images

Western blot analysis of extracts from A431 cells, untreated or treated with EGF, 100ng/ml, using Phospho-mouse ERBB2 Antibody (Y1140)(upper) or Beta-actin (lower).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.