

# OASL Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5164

## Product Information

---

Application	IHC-P, WB
Primary Accession	<a href="#">Q15646</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59226
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	8638
Antigen Region	484-514
Other Names	OASL; TRIP14; 2'-5'-oligoadenylate synthase-like protein; 2'-5'-OAS-related protein; 59 kDa 2'-5'-oligoadenylate synthase-like protein; Thyroid receptor-interacting protein 14; p59 OASL
Dilution	IHC-P~~1:100~500 WB~~1:1000
Target/Specificity	This OASL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 484-514 amino acids from the C-terminal region of human OASL.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	OASL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	OASL
Synonyms	TRIP14

<b>Function</b>	Does not have 2'-5'-OAS activity, but can bind double- stranded RNA. Displays antiviral activity against encephalomyocarditis virus (EMCV) and hepatitis C virus (HCV) via an alternative antiviral pathway independent of RNase L.
<b>Cellular Location</b>	[Isoform p56]: Nucleus, nucleolus. Cytoplasm.
<b>Tissue Location</b>	Expressed in most tissues, with the highest levels in primary blood Leukocytes and other hematopoietic system tissues, colon, stomach and to some extent in testis


## Background

2-prime,5-prime oligoadenylates (2-5As) bind to and activate RNase L, leading to degradation of RNA and inhibition of protein synthesis. 2-5As are produced by 2-5A synthetases (OASs), a highly-conserved family of interferon-induced enzymes. The predicted 514-amino acid human p59OASL (2-5A synthetases-like) protein shares a highly conserved N-terminal domain with other OASs. The C-terminal portion of p59OASL contains 2 ubiquitin-like domains. p59OASL is expressed in most tissues, with the highest levels in hematopoietic tissues, colon, and stomach.


## References

Hovnanian, A., et al., Genomics 56(3):362-363 (1999).  
 Rebouillat, D., et al., Eur. J. Biochem. 257(2):319-330 (1998).  
 Hartmann, R., et al., Nucleic Acids Res. 26(18):4121-4128 (1998).  
 Lee, J.W., et al., Mol. Endocrinol. 9(2):243-254 (1995).  
 Mackay, V., et al., J. Biol. Chem. 251(12):3716-3719 (1976).

## Images



Western blot analysis of lysates from U-87 MG, PC-3, HT-29 cell line (from left to right), using OASL Antibody (T499)(Cat. #AW5164). AW5164 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.