

# BAIAP2 Antibody (C-term)

Mouse Monoclonal Antibody (Mab)

Catalog # AW5188

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q9UQB8</a>
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Calculated MW	60868
Isotype	IgG2b
Antigen Source	Human

## Additional Information

---

Gene ID	10458
Antigen Region	1-290
Other Names	BAIAP2; Brain-specific angiogenesis inhibitor 1-associated protein 2; Brain-specific angiogenesis inhibitor 1-associated protein 2; Fas ligand-associated factor 3; Brain-specific angiogenesis inhibitor 1-associated protein 2; Insulin receptor substrate p53/p58; Brain-specific angiogenesis inhibitor 1-associated protein 2; Insulin receptor substrate protein of 53 kDa
Dilution	WB~~1:1000
Target/Specificity	Purified His-tagged BAIAP2 protein was used to produced this monoclonal antibody.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BAIAP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	BAIAP2
Function	Adapter protein that links membrane-bound small G-proteins to cytoplasmic

effector proteins. Necessary for CDC42-mediated reorganization of the actin cytoskeleton and for RAC1-mediated membrane ruffling. Involved in the regulation of the actin cytoskeleton by WASF family members and the Arp2/3 complex. Plays a role in neurite growth. Acts synergetically with ENAH to promote filipodia formation. Plays a role in the reorganization of the actin cytoskeleton in response to bacterial infection. Participates in actin bundling when associated with EPS8, promoting filopodial protrusions.

#### Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Cell projection, filopodium. Cell projection, ruffle. Cytoplasm, cytoskeleton. Note=Detected throughout the cytoplasm in the absence of specific binding partners. Detected in filopodia and close to membrane ruffles. Recruited to actin pedestals that are formed upon infection by bacteria at bacterial attachment sites

#### Tissue Location

Isoform 1 and isoform 4 are expressed almost exclusively in brain. Isoform 4 is barely detectable in placenta, prostate and testis. A short isoform is ubiquitous, with the highest expression in liver, prostate, testis and placenta


## Background

Adapter protein that links membrane-bound small G-proteins to cytoplasmic effector proteins. Necessary for CDC42-mediated reorganization of the actin cytoskeleton and for RAC1-mediated membrane ruffling. Involved in the regulation of the actin cytoskeleton by WASF family members and the Arp2/3 complex. Plays a role in neurite growth. Acts synergetically with ENAH to promote filipodia formation. Plays a role in the reorganization of the actin cytoskeleton in response to bacterial infection.

## References

Oda K., et al. Cytogenet. Cell Genet. 84:75-82(1999).  
Okamura-Oho Y., et al. Hum. Mol. Genet. 8:947-957(1999).  
Miyahara A., et al. J. Hum. Genet. 48:410-414(2003).  
Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.  
Hachiya T., et al. Submitted (SEP-1996) to the EMBL/GenBank/DDBJ databases.

## Images



Western blot analysis of lysates from K562 cell line, mouse brain tissue, U-251 MG cell line (from left to right), using BAIAP2 Antibody (C-term)(Cat. #AW5188). AW5188 was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.