

R Cebpd Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5199

Product Information

Application	WB
Primary Accession	Q03484
Reactivity	Mouse, Rat, Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28600
Isotype	Rabbit IgG
Antigen Source	Rat

Additional Information

Gene ID	25695
Antigen Region	161-189
Other Names	Cebpd;Celf; CCAAT/enhancer-binding protein delta; CCAAT/enhancer-binding protein delta; Transcription factor CELF
Dilution	WB~~1:1000
Target/Specificity	This Rat Cebpd antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 161-189 amino acids from the Central region of rat Cebpd.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	R Cebpd Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Cebpd
Synonyms	Celf
Function	Transcription activator that recognizes two different DNA motifs: the CCAAT

homology common to many promoters and the enhanced core homology common to many enhancers (PubMed:[1714459](#)). Important transcription factor regulating the expression of genes involved in immune and inflammatory responses. Transcriptional activator that enhances IL6 transcription alone and as heterodimer with CEBPB (By similarity).

Cellular Location

Nucleus.

Tissue Location

Ubiquitously expressed.


Background

C/EBP is a DNA-binding protein that recognizes two different motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers. Important transcriptional activator in the regulation of genes involved in immune and inflammatory responses, may play an important role in the regulation of the several genes associated with activation and/or differentiation of macrophages (By similarity).

References

Kageyama R., et al. J. Biol. Chem. 266:15525-15531(1991).

Images



Western blot analysis of lysates from A549, NCI-H460, mouse NIH/3T3 cell line, rat lung, rat testis tissue, Hela cell line (from left to right), using (RAT) Cebpd Antibody (Center) (Cat. #AW5199). AW5199 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.