

# TUFM Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5218

## Product Information

---

<b>Application</b>	IF, WB
<b>Primary Accession</b>	<a href="#">P49411</a>
<b>Other Accession</b>	<a href="#">P85834</a> , <a href="#">Q8BFR5</a> , <a href="#">P49410</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Predicted</b>	Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Calculated MW</b>	49875
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	7284
<b>Antigen Region</b>	112-147
<b>Other Names</b>	Elongation factor Tu, mitochondrial, EF-Tu, P43, TUFM
<b>Dilution</b>	IF~~1:25 WB~~1:1000
<b>Target/Specificity</b>	This TUFM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 112-147 amino acids from the N-terminal region of human TUFM.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TUFM Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TUFM
<b>Function</b>	GTP hydrolase that promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis. Participates in

mitochondrial translation (By similarity). Also plays a role in the regulation of autophagy and innate immunity (PubMed:[22749352](#), PubMed:[28407488](#)). Recruits ATG5-ATG12 and NLRX1 at mitochondria and serves as a checkpoint of the RIGI-MAVS pathway (PubMed:[28407488](#)). In turn, inhibits RLR-mediated type I interferon while promoting autophagy (PubMed:[22749352](#)).

#### Cellular Location

Mitochondrion matrix

## Background

This protein promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis.

## References

Woriak V.L., et al. *Biochim. Biophys. Acta* 1264:347-356(1995).


Wells J., et al. *FEBS Lett.* 358:119-125(1995).

Ling M., et al. *Gene* 197:325-336(1997).

Martin J., et al. *Nature* 432:988-994(2004).

Dunn M.J., et al. Submitted (MAR-1996) to UniProtKB.

## Images



All lanes : Anti-TUFM Antibody (N-term) at 1:2000 dilution  
Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: Mouse brain tissue lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Rat brain tissue lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.