

TUFM Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5218

Product Information

Application	IF, WB
Primary Accession	P49411
Other Accession	P85834 , Q8BFR5 , P49410
Reactivity	Human, Mouse, Rat
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Calculated MW	49875
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	7284
Antigen Region	112-147
Other Names	Elongation factor Tu, mitochondrial, EF-Tu, P43, TUFM
Dilution	IF~1:25 WB~1:1000
Target/Specificity	This TUFM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 112-147 amino acids from the N-terminal region of human TUFM.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TUFM Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TUFM
Function	GTP hydrolase that promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis. Also plays a role in the

regulation of autophagy and innate immunity. Recruits ATG5-ATG12 and NLRX1 at mitochondria and serves as a checkpoint of the RIGI-MAVS pathway. In turn, inhibits RLR-mediated type I interferon while promoting autophagy.

Cellular Location

Mitochondrion.

Background

This protein promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis.

References

Woriax V.L.,et al.Biochim. Biophys. Acta 1264:347-356(1995).


Wells J.,et al.FEBS Lett. 358:119-125(1995).

Ling M.,et al.Gene 197:325-336(1997).

Martin J.,et al.Nature 432:988-994(2004).

Dunn M.J.,et al.Submitted (MAR-1996) to UniProtKB.

Images



All lanes : Anti-TUFM Antibody (N-term) at 1:2000 dilution
Lane 1: Hela whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: Mouse brain tissue lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Rat brain tissue lysate
Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.