10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PRDX2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5227

Product Information

Application	IF, IHC-P, WB
Primary Accession	P32119
Other Accession	P35704, Q61171, Q2PFZ3, Q8K3U7, Q9BGI3
Reactivity	Mouse, Rat, Human
Predicted	Hamster, Monkey, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21892
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	7001
Antigen Region	169-198
Other Names	PRDX2; NKEFB; TDPX1; Peroxiredoxin-2; Natural killer cell-enhancing factor B; PRP; Thiol-specific antioxidant protein; Thioredoxin peroxidase 1; Thioredoxin-dependent peroxide reductase 1
Dilution	IF~~1:10~50 IHC-P~~1:100~500 WB~~1:1000
Target/Specificity	This PRDX2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 169-198 amino acids from the C-terminal region of human PRDX2.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PRDX2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PRDX2

Synonyms	NKEFB, TDPX1
Function	Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).
Cellular Location	Cytoplasm.


Background

PRDX2 is a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen peroxide and alkyl hydroperoxides. This protein may play an antioxidant protective role in cells, and may contribute to the antiviral activity of CD8(+) T-cells. This protein may have a proliferative effect and play a role in cancer development or progression. The crystal structure of this protein has been resolved to 2.7 angstroms.


References

Engstrom,K.S., et.al., Mutat. Res. (2009)

Images




Western blot analysis of lysates from human brain,human heart,mouse brain,mouse heart,rat heart tissue lysate (from left to right), using PRDX2 Antibody (C-term)(Cat. #AW5227). AW5227 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.



Formalin-fixed and paraffin-embedded human brain tissue reacted with PRDX2 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Fluorescent confocal image of Hela cell stained with PRDX2 Antibody (C-term)(Cat#AW5227).Hela cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with PRDX2 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic



actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min).PRDX2 immunoreactivity is localized to Cytoplasm significantly.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.