

# NRAS Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5296

## Product Information

---

<b>Application</b>	FC, IHC-P, WB
<b>Primary Accession</b>	<a href="#">P01111</a>
<b>Other Accession</b>	<a href="#">Q2MJK3</a>
<b>Reactivity</b>	Mouse, Rat, Human
<b>Predicted</b>	Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	21229
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	4893
<b>Antigen Region</b>	147-179
<b>Other Names</b>	NRAS; HRAS1; GTPase NRas; Transforming protein N-Ras
<b>Dilution</b>	FC~~1:10~50 IHC-P~~1:100~500 WB~~1:1000
<b>Target/Specificity</b>	This NRAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 147-179 amino acids from the C-terminal region of human NRAS.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	NRAS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NRAS
<b>Synonyms</b>	HRAS1

<b>Function</b>	Ras proteins bind GDP/GTP and possess intrinsic GTPase activity.
<b>Cellular Location</b>	Cell membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor Note=Shuttles between the plasma membrane and the Golgi apparatus


## Background

NRAS is a membrane protein that shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. This protein, which has intrinsic GTPase activity, is activated to a GTP-bound form by a GTPase activating protein and inactivated to a GDP-bound form by a guanine nucleotide-exchange factor. Defects in the gene encoding this protein are a cause of juvenile myelomonocytic leukemia (JMML).


## References

Smalley,K.S., Cancer Res. 68 (14), 5743-5752 (2008)  
 Banerji,U., Mol. Cancer Ther. 7 (4), 737-739 (2008)

## Images




Western blot analysis of lysates from MCF-7,A375 cell line (from left to right), using NRAS Antibody (C-term)(Cat. #AW5296). AW5296 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with NRAS antibody (C-term) , which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

NRAS Antibody (C-term) (Cat. #AW5296) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.