

# TIMM50 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5314

#### **Product Information**

**Application** WB **Primary Accession 03ZC08** Reactivity Human Host Rabbit Clonality polyclonal **Calculated MW** 39646 Isotype Rabbit IgG **Antigen Source HUMAN** 

#### **Additional Information**

**Gene ID** 92609

Antigen Region 25-56

**Other Names** Mitochondrial import inner membrane translocase subunit TIM50, TIMM50,

TIM50

**Dilution** WB~~1:1000

**Target/Specificity** This TIMM50 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 25-56 amino acids from the N-terminal

region of human TIMM50.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** TIMM50 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name TIMM50

Synonyms TIM50

**Function** Essential component of the TIM23 complex, a complex that mediates the

translocation of transit peptide-containing proteins across the mitochondrial inner membrane (PubMed:30190335, PubMed:38828998). Has some phosphatase activity in vitro; however such activity may not be relevant in vivo.

**Cellular Location** Mitochondrion inner membrane; Single-pass membrane protein

**Tissue Location** Widely expressed. Expressed at higher level in brain, kidney and liver (at

protein level)


## **Background**

Essential component of the TIM23 complex, a complex that mediates the translocation of transit peptide-containing proteins across the mitochondrial inner membrane. Has some phosphatase activity in vitro; however such activity may not be relevant in vivo. Isoform 2 may participate in the release of snRNPs and SMN from the Cajal body.

#### References

Guo Y., et al.J. Biol. Chem. 279:24813-24825(2004). Zheng H., et al. Submitted (OCT-2003) to the EMBL/GenBank/DDBJ databases. Grimwood J., et al. Nature 428:529-535(2004). Zhang C., et al. Submitted (FEB-1999) to the EMBL/GenBank/DDBJ databases. Bienvenut W.V., et al. Submitted (JUN-2005) to UniProtKB.

### **Images**



Western blot analysis of lysates from Hela, HepG2, SH-SY5Y cell line (from left to right), using TIMM50 Antibody (N-term)(Cat. #AW5314). AW5314 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.