

# TIMM50 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AW5314

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q3ZCQ8</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	polyclonal
<b>Calculated MW</b>	39646
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	92609
<b>Antigen Region</b>	25-56
<b>Other Names</b>	Mitochondrial import inner membrane translocase subunit TIM50, TIMM50, TIM50
<b>Dilution</b>	WB~~1:1000
<b>Target/Specificity</b>	This TIMM50 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 25-56 amino acids from the N-terminal region of human TIMM50.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TIMM50 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TIMM50
<b>Synonyms</b>	TIM50
<b>Function</b>	Essential component of the TIM23 complex, a complex that mediates the

translocation of transit peptide-containing proteins across the mitochondrial inner membrane (PubMed:[30190335](#), PubMed:[38828998](#)). Has some phosphatase activity in vitro; however such activity may not be relevant in vivo.

**Cellular Location** Mitochondrion inner membrane; Single-pass membrane protein

**Tissue Location** Widely expressed. Expressed at higher level in brain, kidney and liver (at protein level)


## Background

Essential component of the TIM23 complex, a complex that mediates the translocation of transit peptide-containing proteins across the mitochondrial inner membrane. Has some phosphatase activity in vitro; however such activity may not be relevant in vivo. Isoform 2 may participate in the release of snRNPs and SMN from the Cajal body.

## References

Guo Y., et al. *J. Biol. Chem.* 279:24813-24825(2004).  
Zheng H., et al. Submitted (OCT-2003) to the EMBL/GenBank/DDBJ databases.  
Grimwood J., et al. *Nature* 428:529-535(2004).  
Zhang C., et al. Submitted (FEB-1999) to the EMBL/GenBank/DDBJ databases.  
Bienvenut W.V., et al. Submitted (JUN-2005) to UniProtKB.

## Images



Western blot analysis of lysates from HeLa, HepG2, SH-SY5Y cell line (from left to right), using TIMM50 Antibody (N-term)(Cat. #AW5314). AW5314 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.