

LRP12 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5323

Product Information

Application	FC, IF, IHC-P, WB
Primary Accession	Q9Y561
Other Accession	Q8BUJ9 , Q9BE74 , NP_038465.1
Reactivity	Mouse, Human
Predicted	Mouse, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94984
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	29967
Antigen Region	635-662
Other Names	LRP12; ST7; Low-density lipoprotein receptor-related protein 12; Suppressor of tumorigenicity 7 protein
Dilution	FC~~1:10~50 IF~~1:10~50 IHC-P~~1:100~500 WB~~1:1000
Target/Specificity	This LRP12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 635-662 amino acids from the C-terminal region of human LRP12.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LRP12 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LRP12
Synonyms	ST7

Function	Probable receptor, which may be involved in the internalization of lipophilic molecules and/or signal transduction. May act as a tumor suppressor.
Cellular Location	Membrane; Single-pass type I membrane protein. Membrane, coated pit
Tissue Location	Widely expressed in heart, skeletal muscle, brain, lung, placenta and pancreas, but not in tissues consisting of a large number of epithelial cells, such as liver and kidney. Expressed at very low levels in a number of tumor-derived cell lines


Background

This gene encodes a member of the low-density lipoprotein receptor related protein family. The product of this gene is a transmembrane protein that is differentially expressed in many cancer cells. Alternate splicing results in multiple transcript variants.


References

Garnis, C., et al. *Oncogene* 23(14):2582-2586(2004)
 Battle, M.A., et al. *Biochemistry* 42(24):7270-7282(2003)
 Qing, J., et al. *Oncogene* 18(2):335-342(1999)

Images





Western blot analysis of lysates from human heart, human skeletal muscle tissue lysate (from left to right), using LRP12 Antibody (C-term) (Cat. #AW5323). AW5323 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.



LRP12 Antibody (C-term) (Cat. #AW5323) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of LRP12 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Confocal immunofluorescent analysis of LRP12 Antibody (C-term) (Cat. #AW5323) with U-251MG cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



LRP12 Antibody (C-term) (Cat. #AW5323) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.