

# BRAF Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5325

## Product Information

---

Application	FC, WB
Primary Accession	<a href="#">P15056</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	84437
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	673
Antigen Region	4-385
Other Names	BRAF; BRAF1; RAFB1; Serine/threonine-protein kinase B-raf; Proto-oncogene B-Raf; p94; v-Raf murine sarcoma viral oncogene homolog B1
Dilution	FC~~1:10~50 WB~~1:1000
Target/Specificity	This BRAF antibody is generated from rabbits immunized with BRAF recombinant protein.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BRAF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	BRAF ( <a href="#">HGNC:1097</a> )
Synonyms	BRAF1, RAFB1
Function	Protein kinase involved in the transduction of mitogenic signals from the cell membrane to the nucleus (Probable). Phosphorylates MAP2K1, and thereby

activates the MAP kinase signal transduction pathway (PubMed:[21441910](#), PubMed:[29433126](#)). Phosphorylates PFKFB2 (PubMed:[36402789](#)). May play a role in the postsynaptic responses of hippocampal neurons (PubMed:[1508179](#)).

#### Cellular Location

Nucleus. Cytoplasm. Cell membrane. Note=Colocalizes with RGS14 and RAF1 in both the cytoplasm and membranes.

#### Tissue Location

Brain and testis.


## Background

BRAF, a member of the RAF subfamily of Ser/Thr protein kinases, is involved in the transduction of mitogenic signals from the cell membrane to the nucleus. It may play a role in the postsynaptic responses of hippocampal neurons. This cytoplasmic protein is expressed in brain and testis. Defects in BRAF are involved in a wide range of cancers including lung cancer and non-Hodgkin lymphoma (NHL). This protein contains 1 zinc-dependent phorbol-ester and DAG binding domain.


## References

Hingorani, S.R., et al., Cancer Res. 63(17):5198-5202 (2003).  
Lee, J.W., et al., Br. J. Cancer 89(10):1958-1960 (2003).  
Davies, H., et al., Nature 417(6892):949-954 (2002).  
Naoki, K., et al., Cancer Res. 62(23):7001-7003 (2002).  
Stephens, R.M., et al., Mol. Cell. Biol. 12(9):3733-3742 (1992).

## Images



Western blot analysis of lysates from Jurkat, K562 cell line (from left to right), using BRAF Antibody (Cat. #AW5325). AW5325 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20 µg per lane.



BRAF Antibody (Cat. #AW5325) flow cytometric analysis of CEM cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.