

EPCAM Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5357

Product Information

Application	WB
Primary Accession	P16422
Other Accession	Q75QW1 , Q3T0L5
Reactivity	Human
Predicted	Mouse, Rat, Bovine
Host	Rabbit
Clonality	polyclonal
Calculated MW	34932
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	4072
Antigen Region	299-334
Other Names	Epithelial cell adhesion molecule, Ep-CAM, Adenocarcinoma-associated antigen, Cell surface glycoprotein Trop-1, Epithelial cell surface antigen, Epithelial glycoprotein, EGP, Epithelial glycoprotein 314, EGP314, hEGP314, KS 1/4 antigen, KSA, Major gastrointestinal tumor-associated protein GA733-2, T
Dilution	WB~~1:1000
Target/Specificity	This EPCAM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 299-334 amino acids from the C-terminal region of human EPCAM.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EPCAM Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EPCAM
-------------	-------

Synonyms	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO
Function	May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.
Cellular Location	Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction. Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction
Tissue Location	Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.


Background

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

References

Strnad J.,et al.Cancer Res. 49:314-317(1989).
Perez M.S.,et al.J. Immunol. 142:3662-3667(1989).
Simon B.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:2755-2759(1990).
Szala S.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:3542-3546(1990).
Linnenbach A.J.,et al.Mol. Cell. Biol. 13:1507-1515(1993).

Images



All lanes : Anti-EPCAM Antibody (C-term)(AW5357) at 1/1000 dilution Lane 1: A431 whole cell lysates Lane 2: HT-29 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.