

PPP2R1B Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5358

Product Information

Application	WB
Primary Accession	P30154
Other Accession	Q4QQT4 , P54613 , Q7TNP2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	66214
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	5519
Antigen Region	86-120
Other Names	Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A beta isoform, PP2A subunit A isoform PR65-beta, PP2A subunit A isoform R1-beta, PPP2R1B
Dilution	WB~~1:1000
Target/Specificity	This PPP2R1B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 86-120 amino acids from the N-terminal region of human PPP2R1B.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PPP2R1B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPP2R1B
Function	The PR65 subunit of protein phosphatase 2A serves as a scaffolding

molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.


Background

The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.

References

Baysal B.E.,et al.Gene 217:107-116(1998).
Wang S.S.,et al.Science 282:284-287(1998).
Baysal B.E.,et al.Eur. J. Hum. Genet. 9:121-129(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Taylor T.D.,et al.Nature 440:497-500(2006).

Images



All lanes : Anti-PPP2R1B Antibody (N-term)(AW5358) at 1/1000 dilution Lane 1: mouse spleen lysates Lane 2: rat kidney lysates Lane 3: SW480 whole cell lysates Lane 4: human testis lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 62 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.