

CHML Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5373

Product Information

Application	IHC-P, FC, WB
Primary Accession	P26374
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74071
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	1122
Antigen Region	624-656
Other Names	Rab proteins geranylgeranyltransferase component A 2, Choroideremia-like protein, Rab escort protein 2, REP-2, CHML, REP2
Dilution	IHC-P~~1:100~500 FC~~1:25 WB~~1:1000
Target/Specificity	This CHML antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 624-656 amino acids from the C-terminal region of human CHML.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CHML Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CHML
Synonyms	REP2
Function	Substrate-binding subunit (component A) of the Rab

geranylgeranyltransferase (GGTase) complex. Binds unprenylated Rab proteins and presents the substrate peptide to the catalytic component B. The component A is thought to be regenerated by transferring its prenylated Rab back to the donor membrane. Less effective than CHM in supporting prenylation of Rab3 family.

Cellular Location

Cytoplasm, cytosol.


Background

Substrate-binding subunit (component A) of the Rab geranylgeranyltransferase (GGTase) complex. Binds unprenylated Rab proteins and presents the substrate peptide to the catalytic component B. The component A is thought to be regenerated by transferring its prenylated Rab back to the donor membrane. Less effective than CHM in supporting prenylation of Rab3 family.


References

Cremers F.P.M.,et al.Hum. Mol. Genet. 1:71-75(1992).
Kasper G.,et al.Gene 295:27-32(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images






All lanes : Anti-CHML Antibody (C-term) at 1/1000 dilution
Lane 1: HT-1080 whole cell lysates Lane 2: HeLa whole cell lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 74 kDa
Blocking/Dilution buffer: 5% NFDm/TBST.



Flow cytometric analysis of HeLa cells using CHML Antibody (C-term)(green, Cat#AW5373) compared to an isotype control of rabbit IgG(blue). AW5373 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

Immunohistochemical analysis of paraffin-embedded H. stomach section using CHML Antibody (C-term)(Cat#AW5373). AW5373 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded R. cerebellum section using CHML Antibody (C-term)(Cat#AW5373). AW5373 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Immunohistochemical analysis of paraffin-embedded M. cerebellum section using CHML Antibody (C-term)(Cat#AW5373). AW5373 was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.