

# Mouse Dbx1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5418

## Product Information

---

Application	WB
Primary Accession	<a href="#">P52950</a>
Other Accession	<a href="#">Q5NSW5</a> , <a href="#">A5PKG8</a>
Reactivity	Mouse, Rat
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36334
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	13172
Antigen Region	261-295
Other Names	Homeobox protein DBX1, Developing brain homeobox protein 1, Dbx1, Dbx
Dilution	WB~~1:1000
Target/Specificity	This Mouse Dbx1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 261-295 amino acids from the C-terminal region of Mouse Dbx1.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Dbx1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	Dbx1
Synonyms	Dbx

<b>Function</b>	Could have a role in patterning the central nervous system during embryogenesis. Has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons. Regulates the transcription factor profile, neurotransmitter phenotype, intraspinal migratory path and axonal trajectory of V0 neurons, features that differentiate them from an adjacent set of V1 neurons.
<b>Cellular Location</b>	Nucleus.


## Background

Could have a role in patterning the central nervous system during embryogenesis. Has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons. Regulates the transcription factor profile, neurotransmitter phenotype, intraspinal migratory path and axonal trajectory of V0 neurons, features that differentiate them from an adjacent set of V1 neurons.

## References

Lu S.,et al.Mech. Dev. 47:187-195(1994).  
Carninci P.,et al.Science 309:1559-1563(2005).  
Pierani A.,et al.Neuron 29:367-384(2001).

## Images



All lanes : Anti-Dbx1 Antibody (C-term) at 1:1000 dilution  
Lane 1: mouse skeletal muscle lysates Lane 2: rat brain lysates  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
Predicted band size : 36 kDa  
Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.