

# MAOA Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5425

## Product Information

---

Application	WB
Primary Accession	<a href="#">P21397</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59682
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	4128
Antigen Region	465-499
Other Names	Amine oxidase [flavin-containing] A, Monoamine oxidase type A, MAO-A, MAOA
Dilution	WB~~1:1000
Target/Specificity	This MAOA antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 465-499 amino acids from the C-terminal region of human MAOA.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MAOA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	MAOA ( <a href="#">HGNC:6833</a> )
Function	Catalyzes the oxidative deamination of primary and some secondary amine such as neurotransmitters, with concomitant reduction of oxygen to hydrogen

peroxide and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues (PubMed:[18391214](#), PubMed:[20493079](#), PubMed:[24169519](#), PubMed:[8316221](#)). Preferentially oxidizes serotonin (PubMed:[20493079](#), PubMed:[24169519](#)). Also catalyzes the oxidative deamination of kynuramine to 3-(2-aminophenyl)-3-oxopropanal that can spontaneously condense to 4-hydroxyquinoline (By similarity).

#### Cellular Location

Mitochondrion outer membrane {ECO:0000250|UniProtKB:P21396}; Single-pass type IV membrane protein {ECO:0000250|UniProtKB:P21396}; Cytoplasmic side {ECO:0000250|UniProtKB:P21396}

#### Tissue Location

Heart, liver, duodenum, blood vessels and kidney.


## Background

Catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

## References

Hsu Y.-P.P.,et al.J. Neurochem. 51:1321-1324(1988).  
Bach A.W.J.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:4934-4938(1988).  
Chen Z.-Y.,et al.Nucleic Acids Res. 19:4537-4541(1991).  
Grimsby J.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:3637-3641(1991).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).

## Images



All lanes : Anti-MAOA Antibody (C-term) at 1:1000 dilution  
Lane 1: human placenta lysates Lane 2: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 60 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.