

# AVPR2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5433

## Product Information

---

<b>Application</b>	WB, FC
<b>Primary Accession</b>	<a href="#">P30518</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	40279
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	554
<b>Antigen Region</b>	343-377
<b>Other Names</b>	Vasopressin V2 receptor, V2R, AVPR V2, Antidiuretic hormone receptor, Renal-type arginine vasopressin receptor, AVPR2, ADHR, DIR, DIR3, V2R
<b>Dilution</b>	FC~~1:25 WB~~1:1000
<b>Target/Specificity</b>	This AVPR2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 343-377 amino acids from the C-terminal region of human AVPR2.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	AVPR2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	AVPR2
<b>Synonyms</b>	ADHR, DIR, DIR3, V2R

<b>Function</b>	Receptor for arginine vasopressin. The activity of this receptor is mediated by G proteins which activate adenylate cyclase. Involved in renal water reabsorption.
<b>Cellular Location</b>	Cell membrane; Multi-pass membrane protein
<b>Tissue Location</b>	Kidney.


## Background

Receptor for arginine vasopressin. The activity of this receptor is mediated by G proteins which activate adenylate cyclase. Involved in renal water reabsorption.


## References

Seibold A., et al. *Am. J. Hum. Genet.* 51:1078-1083(1992).  
 Birnbaumer M., et al. *Nature* 357:333-335(1992).  
 Wildin R.S., et al. *Am. J. Hum. Genet.* 55:266-277(1994).  
 Fay M.J., et al. *Peptides* 17:477-481(1996).  
 North W.G., et al. *Cancer Res.* 58:1866-1871(1998).

## Images



Overlay histogram showing Jurkat cells stained with AW5433 (green line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody ( , 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (1583138) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.



All lanes : Anti-AVPR2 Antibody (C-term) at 1:1000 dilution  
 Lane 1: HeLa whole cell lysates Lane 2: Jurkat whole cell lysates  
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 40 kDa  
 Blocking/Dilution buffer: 5% NFD/MTBST.