

# PYCARD Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5459

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q9ULZ3</a>
Other Accession	<a href="#">NP_037390.2</a> , <a href="#">NP_660183.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21627
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	29108
Antigen Region	129-158
Other Names	Apoptosis-associated speck-like protein containing a CARD, hASC, Caspase recruitment domain-containing protein 5, PYD and CARD domain-containing protein, Target of methylation-induced silencing 1, PYCARD, ASC, CARD5, TMS1
Dilution	WB~~1:1000
Target/Specificity	This PYCARD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 129-158 amino acids from the C-terminal region of human PYCARD.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PYCARD Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	PYCARD {ECO:0000303 Ref.4, ECO:0000312 HGNC:HGNC:16608}
------	---

## Function

Functions as a key mediator in apoptosis and inflammation (PubMed:[11103777](#), PubMed:[12646168](#), PubMed:[15030775](#), PubMed:[17349957](#), PubMed:[17599095](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19234215](#), PubMed:[19494289](#), PubMed:[21487011](#), PubMed:[24630722](#), PubMed:[25847972](#), PubMed:[30674671](#), PubMed:[34678144](#), PubMed:[36050480](#)). Promotes caspase- mediated apoptosis involving predominantly caspase-8 and also caspase-9 in a probable cell type-specific manner (PubMed:[11103777](#), PubMed:[12646168](#)). Involved in activation of the mitochondrial apoptotic pathway, promotes caspase-8-dependent proteolytic maturation of BID independently of FADD in certain cell types and also mediates mitochondrial translocation of BAX and activates BAX-dependent apoptosis coupled to activation of caspase-9, -2 and -3 (PubMed:[14730312](#), PubMed:[16964285](#)). Involved in innate immune response by acting as an integral adapter in the assembly of various inflammasomes (NLRP1, NLRP2, NLRP3, NLRP6, AIM2 and probably IFI16) which recruit and activate caspase-1 leading to processing and secretion of pro-inflammatory cytokines (PubMed:[15030775](#), PubMed:[16982856](#), PubMed:[17349957](#), PubMed:[17599095](#), PubMed:[19158675](#), PubMed:[19158676](#), PubMed:[19234215](#), PubMed:[21487011](#), PubMed:[23530044](#), PubMed:[24630722](#), PubMed:[25847972](#), PubMed:[29440442](#), PubMed:[30674671](#), PubMed:[33980849](#), PubMed:[34678144](#), PubMed:[34706239](#)). Caspase-1-dependent inflammation leads to macrophage pyroptosis, a form of cell death (PubMed:[24630722](#)). The function as activating adapter in different types of inflammasomes is mediated by the pyrin and CARD domains and their homotypic interactions (PubMed:[14499617](#), PubMed:[19234215](#), PubMed:[24630722](#)). Clustered PYCARD nucleates the formation of caspase-1 filaments through the interaction of their respective CARD domains, acting as a platform for of caspase-1 polymerization (PubMed:[24630722](#)). In the NLRP1 and NLRC4 inflammasomes seems not be required but facilitates the processing of procaspase-1 (PubMed:[17349957](#)). In cooperation with NOD2 involved in an inflammasome activated by bacterial muramyl dipeptide leading to caspase-1 activation (PubMed:[16964285](#)). May be involved in RIGI-triggered pro-inflammatory responses and inflammasome activation (PubMed:[19915568](#)). In collaboration with AIM2 which detects cytosolic double-stranded DNA may also be involved in a caspase-1-independent cell death that involves caspase-8 (PubMed:[19158675](#), PubMed:[19158676](#)). In adaptive immunity may be involved in maturation of dendritic cells to stimulate T-cell immunity and in cytoskeletal rearrangements coupled to chemotaxis and antigen uptake may be involved in post-transcriptional regulation of the guanine nucleotide exchange factor DOCK2; the latter function is proposed to involve the nuclear form (PubMed:[22732093](#)). Also involved in transcriptional activation of cytokines and chemokines independent of the inflammasome; this function may involve AP-1, NF-kappa-B, MAPK and caspase-8 signaling pathways (PubMed:[12486103](#), PubMed:[16585594](#)). For regulation of NF-kappa-B activating and inhibiting functions have been reported (PubMed:[12486103](#)). Modulates NF-kappa-B induction at the level of the IKK complex by inhibiting kinase activity of CHUK and IKBK (PubMed:[12486103](#), PubMed:[16585594](#)). Proposed to compete with RIPK2 for association with CASP1 thereby down-regulating CASP1-mediated RIPK2- dependent NF-kappa-B activation and activating interleukin-1 beta processing (PubMed:[16585594](#)). Modulates host resistance to DNA virus infection, probably by inducing the cleavage of and inactivating CGAS in presence of cytoplasmic double-stranded DNA (PubMed:[28314590](#)).

## Cellular Location

Cytoplasm. Inflammasome. Endoplasmic reticulum. Mitochondrion. Nucleus  
Note=Upstream of caspase activation, a redistribution from the cytoplasm to the aggregates occurs. These appear as hollow, perinuclear spherical, ball-like structures (PubMed:[11103777](#), PubMed:[12191486](#), PubMed:[15030775](#)). Upon

NLRP3 inflammasome activation redistributes to the perinuclear space localizing to endoplasmic reticulum and mitochondria (PubMed:12191486, PubMed:15030775). Localized primarily to the nucleus in resting monocytes/macrophages and rapidly redistributed to the cytoplasm upon pathogen infection (PubMed:19234215). Localized to large cytoplasmic aggregate appearing as a speck containing AIM2, PYCARD, CASP8 and bacterial DNA after infection with *Francisella tularensis* (By similarity). {ECO:0000250|UniProtKB:Q9EPB4, ECO:0000269|PubMed:11103777, ECO:0000269|PubMed:12191486, ECO:0000269|PubMed:15030775, ECO:0000269|PubMed:19234215}

### Tissue Location

Widely expressed at low levels. Detected in peripheral blood leukocytes, lung, small intestine, spleen, thymus, colon and at lower levels in placenta, liver and kidney. Very low expression in skeletal muscle, heart and brain. Expressed in lung epithelial cells (at protein level) (PubMed:23229815). Detected in the leukemia cell lines HL-60 and U-937, but not in Jurkat T-cell lymphoma and Daudi Burkitt's lymphoma. Detected in the melanoma cell line WM35, but not in WM793. Not detected in HeLa cervical carcinoma cells and MOLT-4 lymphocytic leukemia cells.


## Background

This gene encodes an adaptor protein that is composed of two protein-protein interaction domains: a N-terminal PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domain (CARD). The PYD and CARD domains are members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the inflammatory and apoptotic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript variants encoding different isoforms have been found for this gene.

## References

Grau, E., et al. J. Cancer Res. Clin. Oncol. 136(9):1415-1421(2010)  
 Motani, K., et al. Cancer Sci. 101(8):1822-1827(2010)  
 Mishra, B.B., et al. Cell. Microbiol. 12(8):1046-1063(2010)  
 Cheng, J., et al. J. Cell. Physiol. 222(3):738-747(2010)  
 McElvania Tekippe, E., et al. PLoS ONE 5 (8), E12320 (2010) :

## Images



All lanes : Anti-PYCARD Antibody (C-term) at 1:1000 dilution  
 Lane 1: THP-1 whole cell lysates  
 Lane 2: MCF-7 whole cell lysates  
 Lysates/proteins at 20 µg per lane.  
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution  
 Predicted band size : 22 kDa  
 Blocking/Dilution buffer: 5% NFDM/TBST.

## Citations

- [Total saponin of Dioscorea collettii attenuates MSU crystal -induced inflammation via inhibiting the activation of the NALP3 inflammasome and caspase -1 in THP -1 macrophages](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.