

UQCRQ Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5464

Product Information

Application	WB
Primary Accession	O14949
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	9906
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	27089
Antigen Region	13-41
Other Names	Cytochrome b-c1 complex subunit 8, Complex III subunit 8, Complex III subunit VIII, Ubiquinol-cytochrome c reductase complex 95 kDa protein, Ubiquinol-cytochrome c reductase complex ubiquinone-binding protein QP-C, UQCRQ
Dilution	WB~~1:1000
Target/Specificity	This UQCRQ antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 13-41 amino acids from the N-terminal region of human UQCRQ.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UQCRQ Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UQCRQ
-------------	-------

Function

Component of the ubiquinol-cytochrome c oxidoreductase, a multisubunit transmembrane complex that is part of the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. The cytochrome b-c1 complex catalyzes electron transfer from ubiquinol to cytochrome c, linking this redox reaction to translocation of protons across the mitochondrial inner membrane, with protons being carried across the membrane as hydrogens on the quinol. In the process called Q cycle, 2 protons are consumed from the matrix, 4 protons are released into the intermembrane space and 2 electrons are passed to cytochrome c.

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P08525};
Single-pass membrane protein {ECO:0000250|UniProtKB:P08525}


Background

This is a component of the ubiquinol-cytochrome c reductase complex (complex III or cytochrome b-c1 complex), which is part of the mitochondrial respiratory chain. This subunit, together with cytochrome b, binds to ubiquinone.

References

Fujiwara T.,et al.Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.
Schaegger H.,et al.Methods Enzymol. 260:82-96(1995).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).
Barel O.,et al.Am. J. Hum. Genet. 82:1211-1216(2008).

Images



All lanes : Anti-UQCRCQ Antibody (N-term) at 1:1000 dilution
Lane 1: human liver lysates
Lane 2: Hela whole cell lysates
Lane 3: MCF-7 whole cell lysates
Lane 4: NCI-H460 whole cell lysates
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 10 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.