

# CYP3A4 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5476

## Product Information

---

Application	FC, IHC-P, WB
Primary Accession	<a href="#">P08684</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57343
Isotype	Rabbit IgG
Antigen Source	HUMAN

## Additional Information

---

Gene ID	1576
Antigen Region	228-255
Other Names	Cytochrome P450 3A4, 11413-, 8-cineole 2-exo-monooxygenase, Albendazole monooxygenase, Albendazole sulfoxidase, CYP3A3, CYP3A4, Cytochrome P450 3A3, Cytochrome P450 H1p, Cytochrome P450 NF-25, Cytochrome P450-PCN1, Nifedipine oxidase, Quinine 3-monooxygenase, Taurochenodeoxycholate 6-alpha-hydroxylase, CYP3A4, CYP3A3
Dilution	FC~~1:10~50 IHC-P~~1:100~500 WB~~1:1000
Target/Specificity	This CYP3A4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 228-255 amino acids from the Central region of human CYP3A4.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CYP3A4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	CYP3A4 {ECO:0000303 PubMed:11470997, ECO:0000312 HGNC:HGNC:2637}
------	---

<b>Function</b>	<p>A cytochrome P450 monooxygenase involved in the metabolism of sterols, steroid hormones, retinoids and fatty acids (PubMed:<a href="#">10681376</a>, PubMed:<a href="#">11093772</a>, PubMed:<a href="#">11555828</a>, PubMed:<a href="#">12865317</a>, PubMed:<a href="#">14559847</a>, PubMed:<a href="#">15373842</a>, PubMed:<a href="#">15764715</a>, PubMed:<a href="#">19965576</a>, PubMed:<a href="#">20702771</a>, PubMed:<a href="#">21490593</a>, PubMed:<a href="#">21576599</a>). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase). Catalyzes the hydroxylation of carbon-hydrogen bonds (PubMed:<a href="#">12865317</a>, PubMed:<a href="#">14559847</a>, PubMed:<a href="#">15373842</a>, PubMed:<a href="#">15764715</a>, PubMed:<a href="#">21490593</a>, PubMed:<a href="#">21576599</a>, PubMed:<a href="#">2732228</a>). Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta- estradiol (E2), namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C-16 position (PubMed:<a href="#">11555828</a>, PubMed:<a href="#">12865317</a>, PubMed:<a href="#">14559847</a>). Plays a role in the metabolism of androgens, particularly in oxidative deactivation of testosterone (PubMed:<a href="#">15373842</a>, PubMed:<a href="#">15764715</a>, PubMed:<a href="#">22773874</a>, PubMed:<a href="#">2732228</a>). Metabolizes testosterone to less biologically active 2beta- and 6beta- hydroxytestosterones (PubMed:<a href="#">15373842</a>, PubMed:<a href="#">15764715</a>, PubMed:<a href="#">2732228</a>). Contributes to the formation of hydroxycholesterols (oxysterols), particularly A-ring hydroxylated cholesterol at the C- 4beta position, and side chain hydroxylated cholesterol at the C-25 position, likely contributing to cholesterol degradation and bile acid biosynthesis (PubMed:<a href="#">21576599</a>). Catalyzes bisallylic hydroxylation of polyunsaturated fatty acids (PUFA) (PubMed:<a href="#">9435160</a>). Catalyzes the epoxidation of double bonds of PUFA with a preference for the last double bond (PubMed:<a href="#">19965576</a>). Metabolizes endocannabinoid arachidonylethanolamide (anandamide) to 8,9-, 11,12-, and 14,15- epoxyeicosatrienoic acid ethanolamides (EpETE-EAs), potentially modulating endocannabinoid system signaling (PubMed:<a href="#">20702771</a>). Plays a role in the metabolism of retinoids. Displays high catalytic activity for oxidation of all-trans-retinol to all-trans-retinal, a rate- limiting step for the biosynthesis of all-trans-retinoic acid (atRA) (PubMed:<a href="#">10681376</a>). Further metabolizes atRA toward 4-hydroxyretinoate and may play a role in hepatic atRA clearance (PubMed:<a href="#">11093772</a>). Responsible for oxidative metabolism of xenobiotics. Acts as a 2-exo- monooxygenase for plant lipid 1,8-cineole (eucalyptol) (PubMed:<a href="#">11159812</a>). Metabolizes the majority of the administered drugs. Catalyzes sulfoxidation of the anthelmintics albendazole and fenbendazole (PubMed:<a href="#">10759686</a>). Hydroxylates antimalarial drug quinine (PubMed:<a href="#">8968357</a>). Acts as a 1,4-cineole 2-exo-monooxygenase (PubMed:<a href="#">11695850</a>). Also involved in vitamin D catabolism and calcium homeostasis. Catalyzes the inactivation of the active hormone calcitriol (1-alpha,25-dihydroxyvitamin D(3)) (PubMed:<a href="#">29461981</a>).</p>
<b>Cellular Location</b>	<p>Endoplasmic reticulum membrane; Single-pass membrane protein.  Microsome membrane; Single-pass membrane protein</p>
<b>Tissue Location</b>	<p>Expressed in prostate and liver. According to some authors, it is not expressed in brain (PubMed:19094056). According to others, weak levels of expression are measured in some brain locations (PubMed:18545703, PubMed:19359404). Also expressed in epithelium of the small intestine and large intestine, bile duct, nasal mucosa, kidney, adrenal cortex, epithelium of the gastric mucosa with intestinal metaplasia, gallbladder, intercalated ducts of the pancreas, chief cells of the parathyroid and the corpus luteum of the ovary (at protein level).</p>

## Background


CYP3A4, is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are

monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs which are used today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens.


## References

Sandanaraj,E., Clin. Cancer Res. 14 (21), 7116-7126 (2008)  
 Nelson,D.R., Pharmacogenetics 14 (1), 1-18 (2004)  
 Inoue,K., Jpn. J. Hum. Genet. 37 (2), 133-138 (1992)


## Images



All lanes : Anti-CYP3A4 Antibody (Center) at 1:1000 dilution Lane 1: HepG2 whole cell lysates Lane 2: A549 whole cell lysates Lane 3: HT-29 whole cell lysates Lane 4: human liver lysates Lane 5: human kidney lysates Lane 6: human ovary lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



CYP3A4 Antibody (Center) (Cat. #AW5476) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CYP3A4 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



CYP3A4 Antibody (Center) (Cat. #AW5476) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.