

GLS Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5478

Product Information

Application IF, IHC-P, WB **Primary Accession** 094925

Other Accession P13264, D3Z7P3
Reactivity Human, Mouse

Predicted Rat
Host Rabbit
Clonality Polyclonal
Calculated MW 73461
Isotype Rabbit IgG
Antigen Source HUMAN

Additional Information

Gene ID 2744

Antigen Region 516-545

Other Names Glutaminase kidney isoform, mitochondrial, GLS, K-glutaminase, L-glutamine

amidohydrolase, GLS, GLS1, KIAA0838

Dilution IF~~1:25 IHC-P~~1:100~500 WB~~1:1000

Target/Specificity This GLS antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 516-545 amino acids from the

C-terminal region of human GLS.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions GLS Antibody (C-term) is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name GLS

Synonyms GLS1, KIAA0838

Function

Catalyzes the first reaction in the primary pathway for the renal catabolism of glutamine. Plays a role in maintaining acid-base homeostasis. Regulates the levels of the neurotransmitter glutamate, the main excitatory neurotransmitter in the brain (PubMed:30239721, PubMed:30575854, PubMed:30970188).

Cellular Location

[Isoform 1]: Mitochondrion {ECO:0000250 | UniProtKB:P13264}. Cytoplasm, cytosol. Note=The 74-kDa cytosolic precursor is translocated into the mitochondria and processed via a 72-kDa intermediate to yield the mature 68- and 65-kDa subunits {ECO:0000250 | UniProtKB:P13264} [Glutaminase kidney isoform, mitochondrial 68 kDa chain]: Mitochondrion matrix {ECO:0000250 | UniProtKB:P13264} Note=Produced by the proteolytic processing of the 74-kDa cytosolic precursor. {ECO:0000250 | UniProtKB:P13264}

Tissue Location

Isoform 1 and isoform 3 are detected in brain cortex. Isoform 3 is highly expressed in astrocytoma, ganglioglioma and ependymoma. Isoform 1 is highly expressed in brain and kidney, but not detected in liver. Isoform 3 is highly expressed in heart and pancreas, detected at lower levels in placenta, lung, pancreas and kidney, but is not detected in liver. Isoform 2 is expressed in cardiac and skeletal muscle.


Background

Sahai (1983) demonstrated phosphate-activated glutaminase (EC 3.5.1.2) in human platelets. It is the major enzyme yielding glutamate from glutamine. Significance of the enzyme derives from its possible implication in behavior disturbances in which glutamate acts as a neurotransmitter(Prusiner, 1981). High heritability of platelet glutaminase was indicated by studies of Sahai and Vogel (1983) [PubMed 6682827] who found an intraclass correlation coefficient of 0.96 for monozygotic twins and 0.53 for dizygotic twins.

References


Swierczynski, I., et.al., Biochim. Biophys. Acta 1157 (1), 55-62 (1993)


Images



All lanes: Anti-GLS Antibody (C-term) at 1:1000 dilution Lane 1: mouse brain lysates Lane 2: human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size: 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Immunohistochemical analysis of paraffin-embedded H.kidney section using GLS Antibody (C-term)(Cat#AW5478). AW5478 was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.





Fluorescent image of Hela cells stained with XAF1 GLS Antibody (C-term)(Cat#AW5478). AW5478 was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.