10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# PPT1 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AW5484

## Product Information

| | |
|---|---|
| **Application** | WB |
| **Primary Accession** | P50897 |
| **Reactivity** | Human |
| **Host** | Mouse |
| **Clonality** | Monoclonal |
| **Calculated MW** | 34193 |
| **Isotype** | IgG1 |
| **Antigen Source** | HUMAN |

## Additional Information

| | |
|---|---|
| **Gene ID** | 5538 |
| **Other Names** | Palmitoyl-protein thioesterase 1, PPT-1, Palmitoyl-protein hydrolase 1, PPT1, PPT |
| **Dilution** | WB~~1:1000 |
| **Target/Specificity** | This PPT1 antibody is generated from a mouse immunized with a full-length recombinant protein from human PPT1. |
| **Format** | Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | PPT1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | PPT1 |
| **Synonyms** | CLN1 {ECO:0000303\|PubMed:19941651}, PPT |
| **Function** | Has thioesterase activity against fatty acid thioesters with 14 -18 carbons, including palmitoyl-CoA, S-palmitoyl-N- acetylcysteamine, and palmitoylated proteins (PubMed:12855696, PubMed:26731412, PubMed:8816748). In contrast to PPT2, PPT1 can hydrolyze palmitoylated proteins and |

palmitoylcysteine (PubMed:12855696).

**Cellular Location** Lysosome. Secreted Golgi apparatus. Endoplasmic reticulum


## Background

Removes thioester-linked fatty acyl groups such as palmitate from modified cysteine residues in proteins or peptides during lysosomal degradation. Prefers acyl chain lengths of 14 to 18 carbons.

## References

Vesa J.,et al.Nature 376:584-587(1995).
Crews C.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:4316-4319(1996).
Schriner J.E.,et al.Genomics 34:317-322(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

## Images




All lanes : Anti-PPT1 Antibody (C-term) at 1:1000 dilution Lane 1: Hela whole cell lysates Lane 2: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.