10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PECI Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5533

Product Information

Application	FC, IHC-P, WB
Primary Accession	O75521
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43585
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	10455
Antigen Region	331-358
Other Names	Enoyl-CoA delta isomerase 2, mitochondrial, DRS-1, Delta(3), delta(2)-enoyl-CoA isomerase, D3, D2-enoyl-CoA isomerase, Diazepam-binding inhibitor-related protein 1, DBI-related protein 1, Dodecenoyl-CoA isomerase, Hepatocellular carcinoma-associated antigen 88, Peroxisomal 3, 2-trans-enoyl-CoA isomerase, pECI, Renal carcinoma antigen NY-REN-1, ECI2, DRS1, HCA88, PECI
Dilution	FC~~1:10~50 IHC-P~~1:100~500 WB~~1:1000
Target/Specificity	This PECI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 331-358 amino acids from the C-terminal region of human PECI.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PECI Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ECI2

Synonyms	DRS1, HCA88, PECI {ECO:0000303|PubMed:10
Function	Auxiliary enzyme in the beta-oxidation of mono- and polyunsaturated fatty acids (PubMed:10419495). Together with the 2,4- dienoyl-CoA reductase (DECR) and the Delta(3,5)-Delta(2,4)-dienoyl-CoA isomerase (ECH), they allow the reentrance of the enoyl-CoA into the beta-oxidation cycle (PubMed:10419495). Able to isomerize both 3-cis (3Z) and 3-trans (3E) double bonds into the 2-trans (2E) form in a range of fatty enoyl-CoA species, with a preference for (3E)-enoyl-CoAs over (3Z)-enoyl-CoAs (PubMed:10419495, PubMed:25515061). Based on its established catalytic mechanism, it is predicted to act also on other fatty acids besides those tested experimentally (Probable).
Cellular Location	[Isoform 1]: Mitochondrion {ECO:0000250|UniProtKB:Q5XIC0}
Tissue Location	Abundant in heart, skeletal muscle and liver. Expressed in CD34(+) T-cells and CD34(+) bone marrow cells


Background

PECI is an auxiliary enzyme that catalyzes an isomerization step required for the beta-oxidation of unsaturated fatty acids.


References

Scanlan,M.J., et.al., Int. J. Cancer 83 (4), 456-464 (1999)
Suk,K., et.al., Biochim. Biophys. Acta 1454 (1), 126-131 (1999)


Images



All lanes : Anti-PECI Antibody (C-term) at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: human heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human skeletal muscle reacted with PECI Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



PECI Antibody (C-term) (Cat. #AW5533) flow cytometric analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.