

# ACTG1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5546

## Product Information

---

<b>Application</b>	IF, FC, WB
<b>Primary Accession</b>	<a href="#">P63261</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	41793
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	71
<b>Antigen Region</b>	188-215
<b>Other Names</b>	Actin, cytoplasmic 2, Gamma-actin, Actin, cytoplasmic 2, N-terminally processed, ACTG1, ACTG
<b>Dilution</b>	IF~~1:10~50 FC~~1:10~50 WB~~1:1000
<b>Target/Specificity</b>	This ACTG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-215 amino acids from the Central region of human ACTG1.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	ACTG1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ACTG1
<b>Synonyms</b>	ACTG
<b>Function</b>	Actins are highly conserved proteins that are involved in various types of cell

motility and are ubiquitously expressed in all eukaryotic cells. May play a role in the repair of noise-induced stereocilia gaps thereby maintains hearing sensitivity following loud noise damage (By similarity).

**Cellular Location** Cytoplasm, cytoskeleton


## Background

Actins are highly conserved proteins that are involved in various types of cell motility, and maintenance of the cytoskeleton. In vertebrates, three main groups of actin isoforms, alpha, beta and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins co-exist in most cell types as components of the cytoskeleton, and as mediators of internal cell motility. Actin, gamma 1, is a cytoplasmic actin found in nonmuscle cells.


## References

de Heer,A.M., Ann. Otol. Rhinol. Laryngol. 118 (5), 382-390 (2009)  
Mouilleron,S., EMBO J. 27 (23), 3198-3208 (2008)  
Liu,P., J Genet Genomics 35 (9), 553-558 (2008)

## Images




All lanes : Anti-ACTG1 Antibody (Center) at 1:1000 dilution  
Lane 1: K562 whole cell lysate Lane 2: MCF-7 whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Flow cytometric analysis of K562 cells using ACTG1 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Confocal immunofluorescent analysis of ACTG1 Antibody (Center)(Cat#AW5546) with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.