

TMED10 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5551

Product Information

Application	WB
Primary Accession	P49755
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24976
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	10972
Antigen Region	156-185
Other Names	Transmembrane emp24 domain-containing protein 10, 21 kDa transmembrane-trafficking protein, S31III125, S31I125, Tmp-21-I, Transmembrane protein Tmp21, p23, p24 family protein delta-1, p24delta1, p24delta, TMED10, TMP21
Dilution	WB~~1:1000
Target/Specificity	This TMED10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-185 amino acids from the C-terminal region of human TMED10.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TMED10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMED10 (HGNC:16998)
Synonyms	TMP21
Function	Cargo receptor involved in protein vesicular trafficking and quality control in the endoplasmic reticulum (ER) and Golgi (PubMed: 10052452 , PubMed: 11726511 , PubMed: 16641999 , PubMed: 17288597 ,

PubMed:[19296914](#), PubMed:[20427317](#), PubMed:[21219331](#), PubMed:[27569046](#)). The p24 protein family is a group of transmembrane proteins that bind coat protein complex I/COPI and coat protein complex II/COPII involved in vesicular trafficking between the membranes (PubMed:[10052452](#)). Acts at the luminal side for incorporation of secretory cargo molecules into transport vesicles and involved in vesicle coat formation at the cytoplasmic side (PubMed:[20427317](#), PubMed:[27569046](#)). Mainly functions in the early secretory pathway and cycles between the ER, ER-Golgi intermediate compartment (ERGIC) and Golgi, mediating cargo transport through COPI and COPII-coated vesicles (PubMed:[10052452](#), PubMed:[10852829](#), PubMed:[12237308](#)). In COPII vesicle-mediated anterograde transport, involved in the transport of GPI-anchored proteins by acting together with TMED2 as their cargo receptor; the function specifically implies SEC24C and SEC24D of the COPII vesicle coat and lipid raft-like microdomains of the ER (PubMed:[20427317](#), PubMed:[27569046](#)). Recognizes GPI anchors structural remodeled in the ER by the GPI inositol-deacylase/PGAP1 and the metallophosphoesterase MPPE1/PGAP5 (By similarity). In COPI vesicle-mediated retrograde transport, involved in the biogenesis of COPI vesicles and vesicle coat recruitment (PubMed:[11726511](#)). Involved in trafficking of amyloid beta A4 protein and soluble APP-beta release (independent from the modulation of gamma-secretase activity) (PubMed:[17288597](#)). Involved in the KDELR2-mediated retrograde transport of the toxin A subunit (CTX-A- K63) together with COPI and the COOH terminus of KDELR2 (By similarity). On Golgi membranes, acts as a primary receptor for ARF1-GDP, a GTP- binding protein involved in COPI-vesicle formation (PubMed:[11726511](#)). Increases coatomer-dependent GTPase-activating activity of ARFGAP2 which mediates the hydrolysis of ARF1-bound GTP and therefore modulates protein trafficking from the Golgi apparatus (PubMed:[19296914](#)). Involved in the exocytic trafficking of G protein-coupled receptors F2LR1/PAR2 (trypsin and trypsin-like enzyme receptor), OPRM1 (opioid receptor) and P2RY4 (UTD and UDP receptor) from the Golgi to the plasma membrane, thus contributing to receptor resensitization (PubMed:[21219331](#)). In addition to its cargo receptor activity, may also act as a protein channel after oligomerization, facilitating the post- translational entry of leaderless cytoplasmic cargo into the ERGIC (PubMed:[32272059](#)). Involved in the translocation into ERGIC, the vesicle entry and the secretion of leaderless cargos (lacking the secretion signal sequence), including the mature form of interleukin 1/IL-1 family members, the alpha-crystallin B chain HSPB5, the carbohydrate-binding proteins galectin-1/LGALS1 and galectin-3/LGALS3, the microtubule-associated protein Tau/MAPT, and the annexin A1/ANXA1; the translocation process is dependent on cargo protein unfolding and enhanced by chaperones HSP90AB1 and HSP90B1/GRP9 (PubMed:[32272059](#)). Could also associates with the presenilin-dependent gamma-secretase complex in order to regulate gamma-cleavages of the amyloid beta A4 protein to yield amyloid-beta 40/Abeta40 (PubMed:[16641999](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Endoplasmic reticulum-Golgi intermediate compartment membrane; Single-pass type I membrane protein. Golgi apparatus membrane; Single-pass type I membrane protein. Golgi apparatus, cis-Golgi network membrane; Single-pass type I membrane protein. Golgi apparatus, trans-Golgi network membrane {ECO:0000250|UniProtKB:Q63584}; Single-pass type I membrane protein. Cytoplasmic vesicle, secretory vesicle membrane; Single-pass type I membrane protein. Cell membrane {ECO:0000250|UniProtKB:Q63584}; Single-pass type I membrane protein. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.


Background

This gene is a member of the EMP24/GP25L/p24 family and encodes a protein with a GOLD domain. This type I membrane protein is localized to the plasma membrane and golgi cisternae and is involved in vesicular protein trafficking. The protein is also a member of a heteromeric secretase complex and regulates the complex's gamma-secretase activity without affecting its epsilon-secretase activity. Mutations in this gene have been associated with early-onset familial Alzheimer's disease. This gene has a pseudogene on chromosome 8.


References

Wang, H., et al. Mol. Biol. Cell 21(8):1398-1408(2010)
 Zhao, J., et al. BMC Med. Genet. 11, 96 (2010) :
 Pardossi-Piquard, R., et al. J. Biol. Chem. 284(42):28634-28641(2009)
 Soranzo, N., et al. PLoS Genet. 5 (4), E1000445 (2009) :
 Liu, S., et al. Eur. J. Neurosci. 28(10):1980-1988(2008)

Images



All lanes : Anti-TMED10 Antibody (C-term) at 1:1000 dilution
 Lane 1: PC-12 whole cell lysate
 Lane 2: NCI-H460 whole cell lysate
 Lane 3: HeLa whole cell lysate
 Lane 4: NIH/3T3 whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-TMED10 Antibody (C-term) at 1:1000 dilution
 Lane 1: PC-12 whole cell lysate
 Lane 2: NCI-H460 whole cell lysate
 Lane 3: HeLa whole cell lysate
 Lane 4: NIH/3T3 whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.