

EWSR1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5593

Product Information

Application	IF, WB
Primary Accession	Q01844
Other Accession	Q61545
Reactivity	Human, Mouse
Predicted	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68478
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	2130
Antigen Region	619-654
Other Names	RNA-binding protein EWS, EWS oncogene, Ewing sarcoma breakpoint region 1 protein, EWSR1, EWS
Dilution	IF~~1:25 WB~~1:2000
Target/Specificity	This EWSR1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 619-654 amino acids from the C-terminal region of human EWSR1.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EWSR1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EWSR1
Synonyms	EWS

Function	Binds to ssRNA containing the consensus sequence 5'-AGGUAA-3' (PubMed: 21256132). Might normally function as a transcriptional repressor (PubMed: 10767297). EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.
Cellular Location	Nucleus. Cytoplasm. Cell membrane. Note=Relocates from cytoplasm to ribosomes upon PTK2B/FAK2 activation
Tissue Location	Ubiquitous.


Background

Might normally function as a transcriptional repressor. EWS-fusion-proteins (EFPS) may play a role in the tumorigenic process. They may disturb gene expression by mimicking, or interfering with the normal function of CTD-POLII within the transcription initiation complex. They may also contribute to an aberrant activation of the fusion protein target genes.


References

Delattre O., et al. Nature 359:162-165(1992).
 Plougastel B., et al. Genomics 18:609-615(1993).
 Zucman-Rossi J., et al. Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.
 Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004).
 Ota T., et al. Nat. Genet. 36:40-45(2004).

Images



All lanes : Anti-EWSR1 Antibody (C-term) at 1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: K562 whole cell lysate Lane 4: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human Cervical epithelial adenocarcinoma cell line) cells labeling EWSR1 with AW5593 at 1/25 dilution, followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/400 dilution (green). Confocal image showing nuclear staining on HeLa cell line. Cytoplasmic actin is detected with Alexa Fluor® 555 conjugated with Phalloidin (OB16636430) at 1/100 dilution (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.