

# HNRNPUL2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5598

## Product Information

---

<b>Application</b>	IHC, WB
<b>Primary Accession</b>	<a href="#">Q1KMD3</a>
<b>Other Accession</b>	<a href="#">Q00PI9</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Dog
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	85105
<b>Isotype</b>	Rabbit IgG
<b>Antigen Source</b>	HUMAN

## Additional Information

---

<b>Gene ID</b>	221092
<b>Antigen Region</b>	317-350
<b>Other Names</b>	Heterogeneous nuclear ribonucleoprotein U-like protein 2, Scaffold-attachment factor A2, SAF-A2, HNRNPUL2, HNRPUL2
<b>Dilution</b>	IHC~~1:100~500 WB~~1:2000
<b>Target/Specificity</b>	This HNRNPUL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 317-350 amino acids from the Central region of human HNRNPUL2.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	HNRNPUL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


---

<b>Name</b>	HNRNPUL2
<b>Synonyms</b>	HNRPUL2


## References

Herrmann F.,et al.Submitted (MAR-2006) to the EMBL/GenBank/DDBJ databases.  
Bechtel S.,et al.BMC Genomics 8:399-399(2007).  
Bienvenut W.V.,et al.Submitted (DEC-2008) to UniProtKB.  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

## Images



All lanes : Anti-HNRNPUL2 Antibody (Center) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AW5598 staining HNRNPUL2 in human stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.