

HNRNPUL2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5598

Product Information

Application	IHC, WB
Primary Accession	Q1KMD3
Other Accession	Q00PI9
Reactivity	Human
Predicted	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85105
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	221092
Antigen Region	317-350
Other Names	Heterogeneous nuclear ribonucleoprotein U-like protein 2, Scaffold-attachment factor A2, SAF-A2, HNRNPUL2, HNRPUL2
Dilution	IHC~~1:100~500 WB~~1:2000
Target/Specificity	This HNRNPUL2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 317-350 amino acids from the Central region of human HNRNPUL2.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HNRNPUL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	HNRNPUL2
Synonyms	HNRPUL2


References

Herrmann F.,et al.Submitted (MAR-2006) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Bienvenut W.V.,et al.Submitted (DEC-2008) to UniProtKB.
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

Images



All lanes : Anti-HNRNPUL2 Antibody (Center) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AW5598 staining HNRNPUL2 in human stomach tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.