

IGF2BP2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5603

Product Information

Application	FC, IHC-P, WB
Primary Accession	Q9Y6M1
Other Accession	Q5SF07 , Q5RB68
Reactivity	Human
Predicted	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66121
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	10644
Antigen Region	530-556
Other Names	Insulin-like growth factor 2 mRNA-binding protein 2, IGF2 mRNA-binding protein 2, IMP-2, Hepatocellular carcinoma autoantigen p62, IGF-II mRNA-binding protein 2, VICKZ family member 2, IGF2BP2, IMP2, VICKZ2
Dilution	FC~~1:10~50 IHC-P~~1:100~500 WB~~1:2000
Target/Specificity	This IGF2BP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 530-556 amino acids from the C-terminal region of human IGF2BP2.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	IGF2BP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IGF2BP2
-------------	---------

Synonyms	IMP2, VICKZ2
Function	RNA-binding factor that recruits target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation (By similarity). Preferentially binds to N6-methyladenosine (m6A)-containing mRNAs and increases their stability (PubMed: 29476152). Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs (PubMed: 9891060). Binding is isoform- specific. Binds to beta-actin/ACTB and MYC transcripts. Increases MYC mRNA stability by binding to the coding region instability determinant (CRD) and binding is enhanced by m6A-modification of the CRD (PubMed: 29476152).
Cellular Location	Nucleus. Cytoplasm. Cytoplasm, P-body. Cytoplasm, Stress granule. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Localizes at the connecting piece and the tail of the spermatozoa. In response to cellular stress, such as oxidative stress, recruited to stress granules
Tissue Location	Expressed in oocytes, granulosa cells of small and growing follicles, Leydig cells, spermatogonia and semen (at protein level). Expressed in testicular cancer (at protein level). Expressed weakly in heart, placenta, skeletal muscle, bone marrow, colon, kidney, salivary glands, testis and pancreas. Detected in fetal liver, fetal ovary, gonocytes and interstitial cells of the testis


Background

This gene encodes a member of the IGF-II mRNA-binding protein (IMP) family. The protein encoded by this gene contains several four KH domains and two RRM domains. It functions by binding to the 5' UTR of the insulin-like growth factor 2 (IGF2) mRNA and regulating IGF2 translation. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.


References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Pechlivanis, S., et al. Arterioscler. Thromb. Vasc. Biol. 30(9):1867-1872(2010)
Henl, M., et al. Diabetes (2010) In press :
Rodriguez, S., et al. Growth Horm. IGF Res. 20(4):310-318(2010)
Voight, B.F., et al. Nat. Genet. 42(7):579-589(2010)


Images



All lanes : Anti-IGF2BP2 Antibody (C-term) at 1:2000 dilution
Lane 1: HepG2 whole cell lysate
Lane 2: HeLa whole cell lysate
Lane 3: A549 whole cell lysate
Lane 4: Raji whole cell lysate
Lane 5: SH-SY5Y whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 66 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



IGF2BP2 antibody(C-term) (Cat. #AW5603)
immunohistochemistry analysis in formalin fixed and
paraffin embedded human spleen tissue followed by
peroxidase conjugation of the secondary antibody and
DAB staining. This data demonstrates the use of the
IGF2BP2 antibody(C-term) for immunohistochemistry.
Clinical relevance has not been evaluated.



IGF2BP2 Antibody (C-term) (Cat. #AW5603) flow
cytometric analysis of HepG2 cells (right histogram)
compared to a negative control cell (left
histogram).FITC-conjugated goat-anti-rabbit secondary
antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.