

NDUFV1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5622

Product Information

| | |
|-------------------|--|
| Application | WB |
| Primary Accession | P49821 |
| Other Accession | P25708 , Q54I90 , Q0MQI5 , Q8HXQ9 , Q91YT0 , Q0MQI6 , Q0MQI4 |
| Reactivity | Human, Mouse |
| Predicted | Human, Dog |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 50817 |
| Isotype | Rabbit IgG |
| Antigen Source | HUMAN |

Additional Information

| | |
|--------------------|---|
| Gene ID | 4723 |
| Antigen Region | 194-226 |
| Other Names | NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial, Complex I-51kD, CI-51kD, NADH dehydrogenase flavoprotein 1, NADH-ubiquinone oxidoreductase 51 kDa subunit, NDUFV1, UQOR1 |
| Dilution | WB~~0.25 |
| Target/Specificity | This NDUFV1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 194-226 amino acids from the Central region of human NDUFV1. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | NDUFV1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|----------|---|
| Name | NDUFV1 (HGNC:7716) |
| Synonyms | UQOR1 |
| Function | Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH |

through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:[28844695](#)). Part of the peripheral arm of the enzyme, where the electrons from NADH are accepted by flavin mononucleotide (FMN) and then passed along a chain of iron-sulfur clusters by electron tunnelling to the final acceptor ubiquinone (PubMed:[28844695](#)). Contains FMN, which is the initial electron acceptor as well as one iron-sulfur cluster (PubMed:[28844695](#)).

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P25708};
Peripheral membrane protein {ECO:0000250|UniProtKB:P25708}; Matrix side {ECO:0000250|UniProtKB:P25708}


Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

References

de Coo R.F.M.,et al.Mamm. Genome 10:49-53(1999).
Schuelke M.,et al.Biochem. Biophys. Res. Commun. 245:599-606(1998).
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-NDUFV1 Antibody (Center) at 1:2000 dilution
Lane 1: HepG2 whole cell lysate
Lane 2: HL-60 whole cell lysate
Lane 3: A431 whole cell lysate
Lane 4: mouse kidney lysate
Lane 5: mouse brain lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.