

ALDOC Antibody (C-term)

Mouse Monoclonal Antibody (Mab)

Catalog # AW5623

Product Information

Application	WB, FC
Primary Accession	P09972
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	39456
Isotype	IgG3
Antigen Source	HUMAN

Additional Information

Gene ID	230
Antigen Region	1-364
Other Names	Fructose-bisphosphate aldolase C, Brain-type aldolase, ALDOC, ALDC
Dilution	WB~~1:2000 FC~~1:25
Target/Specificity	Purified His-tagged ALDOC protein was used to produced this monoclonal antibody.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ALDOC Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	ALDOC
Synonyms	ALDC

References


Rottmann W.H., et al. Biochimie 69:137-145(1987).
Buono P., et al. Nucleic Acids Res. 16:4733-4733(1988).
Buono P., et al. Eur. J. Biochem. 192:805-811(1990).

Yu W., et al. Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.
 Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

Images



Overlay histogram showing HL-60 cells stained with AW5621 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AW5621, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (OJ192088) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG3 ($1\mu\text{g}/1 \times 10^6$ cells) used under the same conditions. Acquisition of >10,000 events was performed.



All lanes : Anti-ALDOC Antibody (C-term) at 1:2000 dilution
 Lane 1: 293T/17 whole cell lysate
 Lane 2: HL-60 whole cell lysate
 Lane 3: rat brain lysate
 Lane 4: mouse brain lysate
 Lysates/proteins at 20 μg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa
 Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.