

MGMT Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AW5643

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, FC |
| Primary Accession | P16455 |
| Other Accession | B4DEE8 |
| Reactivity | Human |
| Predicted | Monkey |
| Host | Mouse |
| Clonality | Monoclonal |
| Calculated MW | 21646 |
| Isotype | IgG1,k |
| Antigen Source | HUMAN |

Additional Information

| | |
|---------------------------|---|
| Gene ID | 4255 |
| Antigen Region | 1-207 |
| Other Names | Methylated-DNA--protein-cysteine methyltransferase, 6-O-methylguanine-DNA methyltransferase, MGMT, O-6-methylguanine-DNA-alkyltransferase, MGMT |
| Dilution | WB~~1:2000 FC~~1:25 |
| Target/Specificity | This MGMT antibody is generated from a mouse immunized with a recombinant protein. |
| Format | Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | MGMT Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|--|
| Name | MGMT |
| Function | Involved in the cellular defense against the biological effects of |

O6-methylguanine (O6-MeG) and O4-methylthymine (O4-MeT) in DNA. Repairs the methylated nucleobase in DNA by stoichiometrically transferring the methyl group to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.

Cellular Location

Nucleus.


Background

Involved in the cellular defense against the biological effects of O6-methylguanine (O6-MeG) in DNA. Repairs alkylated guanine in DNA by stoichiometrically transferring the alkyl group at the O-6 position to a cysteine residue in the enzyme. This is a suicide reaction: the enzyme is irreversibly inactivated.


References

Tano K., et al. Proc. Natl. Acad. Sci. U.S.A. 87:686-690(1990).
Rydberg B., et al. J. Biol. Chem. 265:9563-9569(1990).
Koike G., et al. J. Biol. Chem. 265:14754-14762(1990).
Hayakawa H., et al. J. Mol. Biol. 213:739-747(1990).
Kalnina N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

Images



Overlay histogram showing Jurkat cells stained with AW5643 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AW5643, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (NA168821)) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG (1 µg/1×10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.



All lanes : Anti-MGMT Antibody at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: Molt-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.